

4-16-2020

Seedcorn Maggots Flying in Iowa

Erin Hodgson

Iowa State University, ewh@iastate.edu

Follow this and additional works at: <https://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), and the [Agriculture Commons](#)

Recommended Citation

Hodgson, Erin, "Seedcorn Maggots Flying in Iowa" (2020). *Integrated Crop Management News*. 2623.
<https://lib.dr.iastate.edu/cropnews/2623>

The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.

Seedcorn Maggots Flying in Iowa

Abstract

Seedcorn maggot is a seed and seedling pest of corn and soybean. Plant injury is especially prevalent during cool and wet springs. The larvae, or maggots, feed on germinating corn and soybean seeds or seedlings (Photo 1). Feeding can delay development or kill the plant. Infestations tend to be field-wide instead of in patches like for many other pests. To confirm seedcorn maggot injury, check field areas with stand loss and look for maggots, pupae and damaged seeds (hollowed out seeds or poorly developing seedlings)

Disciplines

Agricultural Science | Agriculture

IOWA STATE UNIVERSITY

Extension and Outreach

Integrated Crop Management

Seedcorn Maggots Flying in Iowa

April 16, 2020

Seedcorn maggot is a seed and seedling pest of corn and soybean. Plant injury is especially prevalent during cool and wet springs. The larvae, or maggots, feed on germinating corn and soybean seeds or seedlings (Photo 1). Feeding can delay development or kill the plant. Infestations tend to be field-wide instead of in patches like for many other pests. To confirm seedcorn maggot injury, check field areas with stand loss and look for maggots, pupae and damaged seeds (hollowed out seeds or poorly developing seedlings).



Photo 1. Typical seedcorn maggot injury in soybean (Photo by Marlin Rice) and corn (Photo by Purdue Extension).

Biology

Seedcorn maggots overwinter in Iowa as pupae in the soil. Adult flies emerge and mate in April and May, then females lay eggs in soil. Maggot densities are higher in soils with high organic matter. Land that is heavily manured may be especially attractive to early-laying females. Recent soil tillage, regardless of residue type, is attractive to egg-laying females.

This fly species has a lower developmental threshold of 39°F and upper threshold of 84°F. Peak adult emergence for the first generation is at 360 accumulated degree days. Based on current accumulated degree days, expect the first generation to be moving around in southern and central Iowa now. There are 4-5 generations of seedcorn maggot per year in our area. Recent freezing temperatures may temporarily slow the development in all areas.




Figure 1. Degree days accumulated (base 39°F) for seedcorn maggot in Iowa (January 1 – April 15, 2020). Map courtesy of Iowa Environmental Mesonet, ISU Department of Agronomy.

Identification

Seedcorn maggots are white, legless and 1/4 inches long with a tapered body (Photo 1). The maggots have a black mouth with hook-like mouthparts to feed. The pupa is brown and looks like a “wheat seed” (Photo 2). The adult fly is grey to brown in color with red eyes. Adult seedcorn maggots are 1/20th inches long and look like a small house fly (Photo 3).



Photo 2. Seedcorn maggot (left) and pupae. Photo by Brian Lang, Iowa State University.



Photo 3. Seedcorn maggot adult is a small, grey fly. Photo by www.ipmimages.org.

Management

There are no rescue treatments for seedcorn maggot. No-till fields are less attractive to egg-laying females. Target planting when soil and moisture conditions are conducive to quick germination and vigorous growth to reduce seed and seedling pest problems. Farmers with persistent seedcorn maggot infestations should consider a later planting date, shallower planting, higher seeding rates, and earlier termination of cover crops (Bessin 2004). Waiting two weeks (or 450 growing degree days) after tillage or manure applications to plant corn or soybean should provide enough time for the seedcorn maggots to complete development and move to another host (Gessell and Calvin 2000).

Insecticidal seed treatments are also an option for persistent seedcorn maggot pressure. If significant stand loss occurs, replanting the field is an option. A replant decision should be based on percent stand loss and cost of additional seed. Corn and soybean resources to aid in replant decisions are available at <https://store.extension.iastate.edu/Product/Soybean-Replant-Decisions>.

References

Bessin, R. 2003. Seedcorn maggots. ENTFACT 309, University of Kentucky Cooperative Extension Service.

Gesell, S., and D. Calvin. 2000. Seed corn maggot as a pest of field corn. Entomological notes, Department of Entomology, Penn State University.

Holm, K. and E. Cullen. 2012. Insect IPM in organic field crops: seedcorn maggot. A3972-01. University of Wisconsin Extension.

Category: Insects and Mites

Links to this article are strongly encouraged, and this article may be republished without further permission if published as written and if credit is given to the author, Integrated Crop Management News, and Iowa State University Extension and Outreach. If this article is to be used in any other manner, permission from the author is required. This article was originally published on April 16, 2020. The information contained within may not be the most current and accurate depending on when it is accessed.

Crops:

Corn Soybean

Tags: seedcorn maggot plant injury seedcorn maggots

Author:



Erin Hodgson *Professor*

Dr. Erin Hodgson started working in the Department of Entomology at Iowa State University in 2009. She is an associate professor with extension and research responsibilities in corn and soybeans. She has a general background in integrated pest management (IPM) for field crops. Dr. Hodgson's curre...