

1-5-2021

Herbicide Programs for Waterhemp Control in Soybean

Prashant Jha

Iowa State University, pjha@iastate.edu

Follow this and additional works at: <https://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), and the [Agriculture Commons](#)

Recommended Citation

Jha, Prashant, "Herbicide Programs for Waterhemp Control in Soybean" (2021). *Integrated Crop Management News*. 2690.

<https://lib.dr.iastate.edu/cropnews/2690>

The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.

Herbicide Programs for Waterhemp Control in Soybean

Abstract

Waterhemp is a dioecious (male and female flowers on separate plants) pigweed with a high genetic diversity. Consequently, this pigweed has a high propensity to adapt to control tactics and has evolved resistance to herbicides from as many as 6 or 7 different herbicide groups (HGs). Out of 75 waterhemp populations collected from corn/soybean fields in Iowa in fall 2019, almost 25% had a four-way multiple resistance to HGs 2 (ALS inhibitors), 5 (atrazine), 9 (glyphosate), and 14 (PPO inhibitors). It is important to use a diverse herbicide program with multiple, effective sites of action to reduce overreliance on any single herbicide and slow further development of herbicide resistance in waterhemp.

Disciplines

Agricultural Science | Agriculture

IOWA STATE UNIVERSITY

Extension and Outreach

Integrated Crop Management

Herbicide Programs for Waterhemp Control in Soybean

January 5, 2021

Waterhemp is a dioecious (male and female flowers on separate plants) pigweed with a high genetic diversity. Consequently, this pigweed has a high propensity to adapt to control tactics and has evolved resistance to herbicides from as many as 6 or 7 different herbicide groups (HGs). Out of 75 waterhemp populations collected from corn/soybean fields in Iowa in fall 2019, almost 25% had a four-way multiple resistance to HGs 2 (ALS inhibitors), 5 (atrazine), 9 (glyphosate), and 14 (PPO inhibitors). It is important to use a diverse herbicide program with multiple, effective sites of action to reduce overreliance on any single herbicide and slow further development of herbicide resistance in waterhemp.

Below are some of the results from recent weed control trials for waterhemp management in soybean conducted by the ISU Weed Science program in Iowa.

Herbicide product	Rate range (oz or fl oz/acre)	Rate evaluated (oz or fl oz/acre)
Verdict	5-10	5
Authority First	6.45-8	5
Authority MTZ	12-20	11
Authority Elite (Autho.Elite)	19-38	28
Enlite	2.8-4.25	2.8
Surveil	3.5-4.2	2.8
Tripzin ZC (Tripz. ZC)	29-58	44
Moccasin MTZ (Mocc.MTZ)	28-56	42
Fierce EZ	6-7.5	6
Fierce MTZ	16-24	16
Panther Pro	12-15	15
Zidua Pro	4.5-6	6

Table 1. Rate range of preemergence (PRE) herbicide products and rates evaluated in ISU field trials in soybean.




Figure 1. Preemergence (PRE) herbicide programs (2 to 3 sites of action, HGs) for waterhemp control in soybean. Dotted red line represents 90% control at 4 weeks after application (WAA).




Figure 2. Two-pass (PRE followed by (fb) POST] herbicide programs for waterhemp control (4 weeks after the POST application) in LibertyLink (LL) or LibertyLink GT27 (LL GT27; treatments with Roundup + Liberty) soybean. Recommended adjuvants were used with all POST herbicides. Dotted red line represents 90% control. Mocc.MTZ,

Autho.Elite, UltBlazer, AnthMax, and CheetaMax refer to Moccasin MTZ, Authority Elite, Ultra Blazer, Anthem Maxx, and Cheetah Max herbicides, respectively.




Figure 3. One pass POST vs. two-pass [PRE followed by (fb) POST] herbicide programs for waterhemp control 4 weeks after application (WAA) of the POST herbicide(s) in Enlist E3 soybean. Recommended adjuvants were used with all POST herbicides. Dotted red line represents 90% control. Mocca. II refers to Moccasin II Plus herbicide.

There is a little difference among PRE products (Figure 1) when appropriate rates are used. Based on Figures 2 and 3, with an effective PRE followed by an effective POST program (two-pass layered residual, with multiple sites-of-action herbicides), there were no significant differences in waterhemp control ($\geq 90\%$; red dotted line) in soybean. For instance, when Verdict PRE at 5 oz/a (70% control) was followed by Liberty (glufosinate) + HG 15 POST, waterhemp control increased to 90% or more. A slight difference in % control observed between the HG 15 herbicides (Outlook, Dual II, vs. Zidua with Roundup + Liberty POST) in Figure 2, could possibly be a rate response.

There is an increased selection pressure for HG 15 used in both corn and soybean. Waterhemp populations with resistance to HG 15 have been reported in the Midwest (not in Iowa yet). Since herbicides from this group are the only ones labeled for use in POST residual programs in soybean, it is recommended to use alternative sites of action including HGs 3, 5, and 14 in conjunction (tank mix/premix) with HG 15 in PRE/burndown programs in the spring (Figures 1, 2, and 3).

2,4-D choline (Enlist One) or glufosinate (Liberty) applied alone POST provided inadequate control (<90%) of waterhemp (Figure 3). In contrast, 2,4-D choline and glufosinate (Interline) applied as a tank-mixture provided superior end-of-season control of waterhemp in Enlist E3 soybean. This strategy (multiple sites of action) is one of the best management practices to delay resistance and preserve the utility of 2,4-D choline and glufosinate for waterhemp control. While these programs evaluated a limited number of available postemergence herbicides, the same principles apply to dicamba-based herbicide programs for Xtend (tolerance to glyphosate and dicamba) and XtendFlex (tolerance to glyphosate, glufosinate, and dicamba) soybeans available for 2021.

Waterhemp has an extended emergence period (May through August in Iowa). Waterhemp grows more rapidly than most weeds prevalent in Iowa soybean fields — up to 1 to 1 ¼ inches per day. Therefore, timing of herbicide applications is critical. Furthermore, a high percentage of waterhemp plants can emerge late in the season (at or after V5-V6 stage of soybean). Plants that emerge or survive under the soybean canopy most likely don't contribute to crop yield reductions; however, those plants can add several thousand seeds (waterhemp being a prolific seed producer) to the soil seed bank. This warrants the need for a season-long management of waterhemp to prevent any seed bank additions using a diversified approach.

Disclaimer: This article is for education purpose only. Mention of a specific product should not be considered as approval, nor should failure to mention a product be considered disapproval. Read the product label before using any herbicide product.

Category: Weeds

Links to this article are strongly encouraged, and this article may be republished without further permission if published as written and if credit is given to the author, Integrated Crop Management News, and Iowa State University Extension and Outreach. If this article is to be used in any other manner, permission from the author is required. This article was originally published on January 5, 2021. The information contained within may not be the most current and accurate depending on when it is accessed.

Crop:

Soybean

Tags: herbicide herbicide resistance weed management

Authors:

Prashant Jha Associate Professor

Prashant Jha is an Associate Professor and Extension Weed Specialist with the Department of Agronomy at ISU. His research program is focused on improved understanding of weed biology and ecology to develop effective, integrated weed management strategies in corn and soybean production systems of ...

Ramawatar Yadav