

2012

Nitrogen sources and sinks in Iowa soils: biogeochemical links between carbon inputs, nitrate leaching, and nitrous oxide emissions

David Mitchell
Iowa State University

Follow this and additional works at: <https://lib.dr.iastate.edu/etd>

 Part of the [Ecology and Evolutionary Biology Commons](#), [Environmental Sciences Commons](#),
and the [Soil Science Commons](#)

Recommended Citation

Mitchell, David, "Nitrogen sources and sinks in Iowa soils: biogeochemical links between carbon inputs, nitrate leaching, and nitrous oxide emissions" (2012). *Graduate Theses and Dissertations*. 12900.
<https://lib.dr.iastate.edu/etd/12900>

This Thesis is brought to you for free and open access by the Iowa State University Capstones, Theses and Dissertations at Iowa State University Digital Repository. It has been accepted for inclusion in Graduate Theses and Dissertations by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

**Nitrogen sources and sinks in Iowa soils: biogeochemical links between
carbon inputs, nitrate leaching, and nitrous oxide emissions**

by

David Christopher Mitchell

A thesis submitted to the graduate faculty

in partial fulfillment of the requirements for the degree of

MASTER OF SCIENCE

Co-majors: Soil Science (Soil Microbiology & Biochemistry);
Ecology and Evolutionary Biology

Program of Study Committee:
Michael J. Castellano, Major Professor
Matthew J. Helmers
Kirsten S. Hofmockel
John E. Sawyer

Iowa State University

Ames, Iowa

2012

Copyright © David Christopher Mitchell, 2012. All Rights Reserved.

TABLE OF CONTENTS

LIST OF TABLES	iv
LIST OF FIGURES	v
ABSTRACT	vi
CHAPTER 1. GENERAL INTRODUCTION	1
Introduction	1
Thesis Organization.....	3
References	4
CHAPTER 2. NITRATE SINKS IN AGRICULTURAL PERENNIAL VEGETATION	
BUFFERS	7
Abstract	7
Keywords	8
Abbreviations	8
Introduction	8
Methods	12
Results	19
Discussion.....	21
Acknowledgements	26
References	26
Tables	31
Figures	32
CHAPTER 3. COVER CROP EFFECTS ON NITROUS OXIDE EMISSIONS	
FROM A MAIZE-BASED CROPPING SYSTEM: INTERACTION OF CARBON	
AND NITROGEN INPUTS	38
Abstract	38
Keywords	39
Introduction	39
Methods	41
Results	46
Discussion.....	48
Acknowledgements	53
References	53
Tables	58

Figures 59

CHAPTER 4. GENERAL CONCLUSION64

ACKNOWLEDGEMENTS66

LIST OF TABLES

Table 2.1. Soil NO ₃ -N concentrations by watershed and vegetation treatment immediately before and immediately following isotope addition	31
Table 2.2. Soil NH ₄ -N and NO ₃ -N concentrations by watershed and vegetation treatment 7 and 137 days after isotope addition	31
Table 3.1. Maize grain yields (at 155 g kg ⁻¹ moisture) at the field site harvested 5 Oct 2011	58
Table 3.2. Analysis of variance of cumulative N ₂ O produced over 10 days in incubating soils	58

LIST OF FIGURES

Figure 2.1. Precipitation at the research site in 2011.....	32
Figure 2.2. Soil particulate organic matter nitrogen (POM-N) concentrations and carbon: nitrogen ratio (POM C:N) by vegetation treatment and time after isotope addition	33
Figure 2.3. Soil mineral-associated organic matter nitrogen (MAOM-N) and carbon (MAOM-C) concentrations by vegetation treatment and time after isotope addition	34
Figure 2.4. Percent recovery of ¹⁵ N tracer in particulate organic matter (POM) and mineral-associated organic matter (MAOM) by vegetation treatment and time after isotope addition	35
Figure 2.5. Percent of ¹⁵ N tracer recovered in aboveground biomass (AGBM), course root biomass (CRBM), fine root biomass (FRBM), particulate organic matter (POM), mineral-associated organic matter (MAOM), and unrecovered in vegetated cores.....	36
Figure 2.6. Comparison of NO ₃ -N concentrations, denitrification enzyme potential, water-filled pore space (WFPS), and dissolved organic carbon (DOC) concentrations in soils from equivalent topographic positions in perennial vegetation buffers and row crops (soybean)	37
Figure 3.1. Mean soil NO ₃ -N concentrations over the period of field gas flux measurements	59
Figure 3.2. Cumulative N ₂ O-N emissions from the field study	60
Figure 3.3. Initial NO ₃ -N concentrations in incubating soils collected from the fertilizer bands of the field site 13-16 June 2011	61
Figure 3.4. Cumulative N ₂ O-N emissions over 10 days from soils collected from the fertilizer bands of the field site and incubated at 0.8 cm ³ cm ⁻³ WFPS, showing the effect of NO ₃ addition at each N fertilizer rate across cover crop and glucose treatments	62
Figure 3.5. Cumulative N ₂ O-N emissions over 10 days from soils collected from the fertilizer bands of the field site and incubated at 0.8 cm ³ cm ⁻³ WFPS, showing the effect of glucose addition at each cover crop × N fertilizer rate combination.....	63

ABSTRACT

Nitrogen (N) lost from agricultural soils in the forms of nitrate (NO_3) and nitrous oxide (N_2O) have become major environmental concerns. Because N cycling is coupled with organic carbon (C) cycling, management practices that influence soil organic C inputs and cycling may affect reactive N losses. Management practices have been proposed to reduce N losses, including perennial vegetation buffers (PVB) and overwintering non-legume cover crops. However, the effects of these practices on N losses depend on the biogeochemical interactions between soil N and C cycling. This thesis presents investigations of the effects of these management practices on NO_3 and N_2O losses from row crop systems in Iowa, USA. In PVBs, soil organic matter and plant biomass acted as sinks for NO_3 inputs. However, denitrification, stimulated by organic C inputs from perennial vegetation, appeared to be the most important NO_3 sink. These results indicate that integration of perennial vegetation into agricultural landscapes can return substantial amounts of N to the atmosphere and decrease watershed NO_3 losses in the long term. The effects of cover crops on N_2O emissions were found to vary with N fertilizer rate, and cover crops increased N_2O emissions at an economical N rate. These results indicate that overwintering cover crops should not be expected to consistently decrease N_2O emissions from agricultural soils, even when they do decrease NO_3 availability for denitrification. In row crop systems with PVBs and cover crops, mineralizable C inputs to soils are a key factor influencing the biogeochemical N transformations that lead to N retention or losses. Considering the interactions between C and N cycling in agricultural soils is necessary to understand and predict the effects of management practices on environmental N losses.

CHAPTER 1. GENERAL INTRODUCTION

Introduction

Nitrogen (N) cycling plays a central role in agricultural production and global change.

Nitrogen fertilizer is a key factor sustaining the high yields of modern agricultural systems (Cassman et al., 2002). However, loss of various forms of reactive N from agricultural ecosystems has had serious environmental consequences (Robertson and Vitousek, 2009).

Globally, human activity has greatly accelerated N cycling, largely due to high inputs of N to agricultural systems; resultant N losses have had great effects on the biosphere (Vitousek et al., 1997; Galloway et al., 2008). Management of N in agricultural systems to sustain productivity while decreasing reactive N losses therefore presents a great challenge (Robertson and Vitousek, 2009).

Maize-based agroecosystems are a major land cover in the United States, accounting for over one-third of cropland (NASS, 2011). Maize-soybean agroecosystems, predominant in the upper Midwestern USA, are characterized by high crop productivity, providing more than one-third of global corn and soybean production (Dobermann and Cassman, 2002). However, substantial N losses also occur from these systems (Vitousek et al., 2009). Notably, nitrate (NO_3) losses from this cropping system have led to contamination of rivers and streams (Hatfield et al., 2009). Nitrate originating from maize-soybean agriculture has been identified as the leading cause of the annual hypoxic zone in the Gulf of Mexico (Alexander et al., 2008; David et al., 2010).

In addition to dissolved NO_3 losses to aquatic ecosystems, emissions of the greenhouse gas nitrous oxide (N_2O) are an environmental concern due to the contribution of N_2O to radiative forcing in the atmosphere (Smith et al., 2007). Agriculture is currently the dominant cause of increasing atmospheric N_2O concentrations (Reay et al., 2012). Though maize-based agricultural systems can also be sources of carbon dioxide and methane, N_2O contributes the most to the total global warming potential of these systems (Robertson et al., 2000; Mosier et al., 2005; Adviento-Borbe et al., 2007). Models of N_2O emissions from agricultural soils have identified the upper Midwestern US as a major source of this greenhouse gas (USEPA, 2010).

There is potential to reduce N losses from agroecosystems by re-coupling nitrogen inputs to biological processes (Drinkwater and Snapp, 2007). Management changes beyond modification of N fertilizer rate and timing are likely necessary to substantially decrease reactive N losses (Hatfield et al., 2009). Since N cycling processes in soil are closely coupled to soil organic carbon (SOC) (Booth et al., 2005), the effectiveness of practices meant to reduce N losses depends in part on the links between N cycling and soil C inputs and cycling (Drinkwater and Snapp, 2007). Among the practices proposed to decrease reactive N losses from croplands are increasing land cover by perennial vegetation (Schulte et al., 2006) and replacing winter fallow periods with cover crops (Thorup-Kristensen et al., 2003).

Thesis Organization

This thesis presents studies of two management practices aimed to reduce reactive N losses from soils in maize-soybean agroecosystems in Iowa, USA. These studies focus on N losses as NO_3 (Chapter 2) and as N_2O (Chapter 3) in relation to management practices proposed to reduce these losses. Both studies investigate organic C inputs and cycling as key factors controlling reactive N losses from these soils. The overall conclusions from these studies are synthesized and summarized in Chapter 4.

The study presented in Chapter 2 investigates the NO_3 sink strength of soils managed in perennial vegetation within row crop landscapes. While these perennial vegetation buffers (PVB) have been shown to decrease NO_3 losses to downstream ecosystems, the effectiveness of this ecosystem service depends on the strength of various biogeochemical sinks for NO_3 entering these soils in subsurface flow. Though previous studies of PVB have focused on plant uptake and denitrification, soil organic matter (SOM) may act as a major NO_3 sink as well, since SOM is the largest sink for N in terrestrial ecosystems (Aber et al., 1998). This study links ecosystem nutrient retention theory (Vitousek and Reiners, 1975) to SOM saturation theory (Six et al., 2002) and applies these principles to a management practice meant to improve environmental quality.

The study presented in Chapter 3 investigates the effect of overwintering non-legume cover crops in a maize-soybean rotation on emissions of N_2O from the soil surface. While studies have proposed that cover crops decrease soil N_2O emissions, this effect is not consistent.

Chapter 3 presents evidence that the inconsistent effects of cover crops on N₂O emissions may be influenced by mineral N availability, as influenced by N fertilizer rate. This study draws from previous work examining the interaction between soil mineralizable C and NO₃ availability as controls on N₂O emissions. This study provides evidence that expectations of cover crops to decrease N₂O emissions from the soil may be confounded by the complex interactive effects of cover crops and N management, as well as climate and other factors, on N₂O emissions.

References

- Aber J, McDowell W, Nadelhoffer K, et al. 1998. Nitrogen saturation in temperate forest ecosystems: hypotheses revisited. *BioScience* 48: 921- 934.
- Adviento-Borbe MAA, Haddix ML, Binder DL, Walters DT, Dobermann A. 2007. Soil greenhouse gas fluxes and global warming potential in four high-yielding maize systems. *Global Change Biology* 13: 1972-1988.
- Alexander RB, Smith RA, Schwarz GE, Boyer EW, Nolan JV, Brakebill JW. 2008. Differences in phosphorus and nitrogen delivery to the Gulf of Mexico from the Mississippi River basin. *Environmental Science and Technology* 42: 822-830.
- Booth MS, Stark JM, Rastetter E. 2005. Controls on nitrogen cycling in terrestrial ecosystems: a synthetic analysis of literature data. *Ecological Monographs* 75: 139-157.
- Cassman KG, Dobermann A, Walters DT. 2002. Agroecosystems, nitrogen-use efficiency, and nitrogen management. *Ambio* 31: 132-140.
- David MB, Drinkwater LE, McIsaac GF. 2010. Sources of nitrate yields in the Mississippi River basin. *Journal of Environmental Quality* 39: 1657-1667.
- Dobermann A, Cassman KG. 2002. Plant nutrient management for enhanced productivity in intensive grain production systems of the United States and Asia. *Plant and Soil* 247: 153-175.

- Drinkwater LE, Snapp SS. 2007. Nutrients in agroecosystems: rethinking the management paradigm. *Advances in Agronomy* 92: 163-186.
- Galloway JN, Townsend AR, Erisman JW, et al. 2008. Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science* 320: 889-892.
- Hatfield JL, McMullen LD, Jones CS. 2009. Nitrate-nitrogen patterns in the Raccoon River Basin related to agricultural practices. *Journal of Soil and Water Conservation* 64: 190-199.
- Mosier AR, Halvorson AD, Reule CA, Liu XJ. 2006. Net global warming potential and greenhouse gas intensity in irrigated cropping systems in northeastern Colorado. *Journal of Environmental Quality* 35: 1584-1598.
- National Agricultural Statistics Service, US Department of Agriculture. 2011. <http://www.nass.usda.gov/index.asp>. Accessed October 2012.
- Reay DS, Davidson EA, Smith KE, Smith P, Melillo JM, Dentener F, Crutzen PJ. 2012. Global agriculture and nitrous oxide emissions. *Nature Climate Change* 2: 410-416.
- Robertson GP, Paul EA, Harwood RR. 2000. Greenhouse gases in intensive agriculture: contributions of individual gases to the radiative forcing of the atmosphere. *Science* 289: 1922-1925.
- Robertson GP, Vitousek PM. 2009. Nitrogen in agriculture: balancing the cost of an essential resource. *Annual Review of Environment and Resources* 34: 97-125.
- Schulte LA, Liebman MZ, Asbjornsen H, Crow TR. 2006. Agroecosystem restoration through strategic integration of perennials. *Journal of Soil and Water Conservation* 61: 165-169.
- Smith P, Martino D, Cai, Z, et al. 2007. Chapter 8: Agriculture. In: Metz B, Davidson OR, Bosch PR, Dave R, Meyer LA (Eds.). *Climate Change 2007: Mitigation. Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK, pp. 497-540.
- Six J, Conant RT, Paul EA, Paustian K. 2002. Stabilization mechanisms of soil organic

- matter: implications for C-saturation of soils. *Plant and Soil* 241: 155-176.
- Thorup-Kristensen K, Magid J, Jensen LS. 2003. Catch crops and green manures as biological tools in nitrogen management in temperate zones. *Advances in Agronomy* 79: 227-302.
- US Environmental Protection Agency. 2011. Inventory of Greenhouse Gas Emissions and Sinks. Washington, DC. Accessed October 2012. Available at <http://www.epa.gov/climatechange/emissions/usinventoryreport.html>
- Vitousek PM, Mooney HA, Lubchenco J, Melillo JM. 1997. Human domination of earth's ecosystems. *Science* 277: 494-499.
- Vitousek PM, Naylor R, Crews T, et al. 2009. Nutrient imbalances in agricultural development. *Science* 324: 1519-1520.
- Vitousek PM, Reiners WA. 1975. Ecosystem succession and nutrient retention: a hypothesis. *BioScience* 25: 376-381.

CHAPTER 2. NITRATE SINKS IN AGRICULTURAL PERENNIAL VEGETATION BUFFERS

A paper to be submitted to the journal *Ecosystems*

David C. Mitchell¹, Michael J. Castellano², Timothy B. Parkin³, Matthew J. Helmers⁴

Abstract

Nitrate (NO_3) leaching from agricultural watersheds is a major cause of pollution in aquatic ecosystems. Integration of perennial vegetation into agricultural watersheds may decrease NO_3 losses to surface and groundwaters. However, the long-term effectiveness of this ecosystem service depends on the relative importance of several NO_3 sinks. Vegetation and labile soil organic matter (SOM) are temporary sinks for NO_3 . In contrast, stable SOM is a long-term, but potentially finite, NO_3 sink, while gaseous loss through denitrification is a permanent sink. We investigated the relative importance of NO_3 sinks in perennial vegetation buffers that were integrated into agricultural watersheds in Iowa, USA, and had been shown to decrease soil solution NO_3 concentrations up to 100%. Using a $^{15}\text{NO}_3$ tracer, we quantified $\text{NO}_3\text{-N}$ recovery in vegetation and SOM sinks after one growing season while comparing $\text{NO}_3\text{-N}$ recovery in SOM with and without growing perennial vegetation. We also compared

¹ Graduate student, Department of Agronomy, ISU. Primary researcher and author of the paper.

² Assistant professor, Department of Agronomy, ISU. Major professor of DC Mitchell.

³ Research Scientist, USDA National Laboratory for Agriculture and Environment. Contributed intellectually to this paper.

⁴ Associate professor, Department of Agricultural and Biosystems Engineering, ISU. Contributed intellectually to this paper.

potential denitrification enzyme activity in soils from paired watersheds with and without perennial vegetation buffers. Assimilation of $\text{NO}_3\text{-N}$ into labile and stable SOM pools was rapid and initially independent of vegetation; however, the presence of vegetation increased retention of $\text{NO}_3\text{-N}$ in SOM by the end of the growing season. Nevertheless, $\text{NO}_3\text{-N}$ recovery in SOM and vegetation accounted for <20% of $\text{NO}_3\text{-N}$ inputs. Denitrification enzyme data indicated that labile C inputs from perennial vegetation increased denitrifier activity in soils under perennial vegetation. Together, these results indicate that denitrification may be a more important sink for NO_3 inputs than vegetation and SOM in perennial buffers. Carbon inputs from perennial vegetation appear to be important for NO_3 retention in stable SOM and loss by denitrification.

Keywords: soil organic matter; nitrogen retention; nitrate leaching; denitrification

Abbreviations: PVB, perennial vegetation buffers; NO_3 , nitrate; SOM, soil organic matter; SOC, soil organic carbon; SON, soil organic nitrogen; AGBM, aboveground biomass; FRBM, fine root biomass; CRBM, course root biomass; TSEN, total salt-extractable nitrogen; SEOC, salt-extractable organic carbon; DOC, dissolved organic carbon; DEA, denitrification enzyme assay

Introduction

Nutrient loss from agricultural land has become a leading source of ground and surface water pollution in the United States (Carpenter et al., 1998). In the Upper Mississippi River Basin,

maize (*Zea mays* L.)-soybean [*Glycine max* L. (Merr.)] croplands are the major source of nitrate (NO₃) contamination of streams which is responsible for the annual hypoxic zone in the Gulf of Mexico (Alexander et al., 2008; David et al., 2010). Contamination of drinking water by agriculturally-derived NO₃ also poses risks to human health (Townsend et al., 2003).

Perennial vegetation buffers (PVB) in agricultural landscapes can reduce NO₃ loads to downstream ecosystems by removing NO₃ from runoff (Hill et al., 1996; Mayer et al., 2007). However, PVB can remove NO₃ from agricultural runoff by various mechanisms, including uptake by vegetation, storage in soil organic matter (SOM), and gaseous loss through denitrification. From the perspective of reducing dissolved NO₃ losses, these processes differ in their long-term effectiveness (Schade and Lewis, 2006). According to nutrient retention theory, ecosystems receiving chronic nitrogen (N) inputs have a limited capacity to retain these inputs in biological and organic matter sinks. After this capacity is reached, N outputs in the form of gaseous and dissolved losses are equivalent to N inputs (Vitousek and Reiners, 1975; Aber et al., 1989). Thus, if plant biomass and SOM are the major sinks for NO₃ inputs in PVB, reductions in watershed NO₃ losses observed after PVB implementation may not continue in the long-term. In contrast, gaseous loss of NO₃ through denitrification represents a more permanent sink for NO₃ inputs (Martin et al., 1999; Mayer et al., 2007).

Denitrification has been found or inferred to be the dominant sink for agriculturally-derived NO₃ in many PVB (Vought et al., 1994; Verchot et al., 1997; Martin et al., 1999), including those in which plant demand has been saturated (Hanson et al., 1994a, 1994b). Availability

of mineralizable soil organic carbon (SOC) beneath perennial vegetation can stimulate denitrification (Schade et al., 2001; Baker and Vervier, 2004). During denitrification, $\text{NO}_3\text{-N}$ is lost from the ecosystem as gaseous N species. Thus, if denitrification is the principal sink for NO_3 inputs to PVB, reductions in NO_3 loads to downstream systems would be expected to continue in the long term.

The magnitude of NO_3 uptake by vegetation varies with type and management, but can be the major factor reducing NO_3 losses in some PVB systems (Hefting et al., 2005). However, plant biomass is necessarily a temporary sink for $\text{NO}_3\text{-N}$; biomass-N does not accumulate indefinitely, but instead ultimately either is mineralized or becomes soil organic N (SON) (Vitousek and Reiners 1975). While N mineralized from plant biomass can be readily lost from the soil, SON deriving from biomass may or may not be retained in the long term. Nevertheless, harvest of aboveground biomass can provide a permanent sink for N inputs to PVB.

Though previous research has focused on denitrification and plant biomass, incorporation of $\text{NO}_3\text{-N}$ into SOM is potentially an important sink for NO_3 inputs into PVB. Soil organic matter is the largest N sink in terrestrial ecosystems (Aber et al., 1998). Furthermore, since inorganic N transformations in soil correlate closely with SOC (Booth et al., 2005), SOM potentially plays a central role in $\text{NO}_3\text{-N}$ removal from agricultural runoff passing through PVB. Nitrate-N can be incorporated into SOM through plant uptake and subsequent litter decomposition or directly through microbial and abiotic mechanisms (Barrett and Burke, 2002; Dell et al., 2005).

While a fraction of SOM is available for microbial mineralization, the remainder is relatively stable in the long term. Particulate organic matter (POM), composed of large compounds derived from plant biomass, is considered to be potentially mineralizable by microorganisms (Six et al., 2002). In contrast, SOM that is physico-chemically bound to silt and clay particles, known as mineral-associated organic matter (MAOM), is considered to be relatively resistant to microbial mineralization (Six et al., 2002; Kleber et al., 2007). The amount of SOM that can be protected from mineralization as MAOM is limited and a function of the silt + clay content of the soil (Feng et al., 2012). After the soil MAOM pool reaches capacity (saturates), additional SOM inputs would not be physico-chemically protected from mineralization (Hassink, 1997; Stewart et al., 2007). In this case, as SOM accumulates and N sinks in MAOM saturate, further N inputs would not be protected from re-mineralization and loss (Castellano et al., 2012). Therefore, if SOM is the principal sink for $\text{NO}_3\text{-N}$ inputs to PVB, decreases in watershed NO_3 losses may not continue in the long term due to saturation of stable SOM and active cycling of N retained in mineralizable SOM.

The objective of this study was to evaluate the $\text{NO}_3\text{-N}$ sink strength of plant biomass, SOM, and denitrification in PVB integrated into an agricultural landscape in Iowa, USA. We used an isotope tracer to quantify the NO_3 sink strengths of plant biomass and SOM. Relatively few previous studies have used this method in agricultural PVB (Matheson et al., 2002; Bedard-Haughn et al., 2004), and to our knowledge, no previous studies have used this method in the Midwestern region of the USA. Preliminary data has shown that, since implementation, SON concentrations in these PVB have increased by 50-200% (Pérez-

Suárez, *unpublished data*), showing accumulation of N inputs in SOM. Based on these preliminary results, we have hypothesized that SOM is the most important sink for NO₃ in this system.

Methods

Field Site and Experimental Design

This study was conducted at an experimental site within the Neal Smith National Wildlife Refuge in Jasper County, IA, USA (41°33' N, 93°16' W). This site was established to evaluate PVB as a conservation practice to mitigate soil erosion and NO₃ loss (Zhou et al., 2010). At this site, three pairs of experimental watersheds were established. All watersheds had been managed as unfertilized brome (*Bromus* sp.) for at least 10 yrs prior to 2006. Watersheds were tilled with a mulch tiller in fall 2006 and spring 2007. The following treatments were begun in 2007: perennial vegetation buffers were planted in July 2007 in the lowest 10% of the area of one watershed in each pair, while the remaining 90% of those watersheds and 100% of the area of the other watershed in each pair was managed as a no-till maize-soybean rotation (soybean in 2007). The vegetation community of PVBs at the time of this study was 90% perennial (Hirsh, 2012). Anhydrous ammonia was applied to maize at 134 and 186 kg N ha⁻¹ in 2008 and 2010, respectively. Biomass (>15 cm in height) has been harvested and removed from the PVB in late October or November annually since 2010. Compared to the equivalent landscape position in watersheds managed entirely as row crops, solution NO₃ concentrations were decreased by PVB establishment in vadose zone (50-

100%) and shallow groundwater (60-80%) (Zhou et al., 2010). Though subsurface flow rates could not be quantified at the site, they can be inferred to be lower in watersheds with PVB since evapotranspiration in PVB is expected to be greater than in row crops. Thus, though $\text{NO}_3\text{-N}$ loads from these watersheds could not be calculated, loads from watersheds with PVB are inferred to be decreased compared to from watersheds entirely in row crops.

The $\text{NO}_3\text{-N}$ sink strengths of PVB plant biomass and SOM were measured in the field using a procedure adapted from Dell et al. (2005). Three pairs of polyvinyl chloride (PVC) cores (25 cm diameter \times 30 cm height, open on both ends) were pushed completely into the soil in each PVB. The pairs of PVC cores were installed at random locations within 5 m of the uphill edge of each PVB; this placement was chosen based on the assumption that N retention processes would be most important close to the interface between cropland and perennial vegetation, as found in similar studies of ecosystems receiving hydrologic N inputs (Lowrance et al., 1992; Haycock and Pinay, 1993). Since Zhou et al. (2010) had shown decreased NO_3 losses in watersheds with PVB compared to watersheds without, cores were not installed in watersheds without PVB. Instead, the focus of this study was to evaluate the strength of sinks within the PVB. Perennial vegetation emerged in all PVC cores after installation.

On 11 May 2011, one PVC core from each pair was randomly selected for vegetation removal. Vegetation in these cores (hereafter “de-vegetated cores”) was cut to ground level. Glyphosate [*N*-(phosphonomethyl)glycine] solution (20% in water) was brushed onto cut vegetation with a paintbrush. The cut above-ground biomass was then placed on the soil

surface within the cores. Perennial plants did not re-emerge in de-vegetated cores for the remainder of the growing season, indicating that herbicide application was effective. Weed seedlings were removed manually from the de-vegetated cores periodically throughout the growing season. Vegetation in the other core from each pair (hereafter “vegetated cores”) was allowed to grow undisturbed throughout the growing season.

Isotope Addition and Initial Soil Data

On 15 June 2011, 1 L of 3.63 mM KNO₃ (54.5 mg NO₃-N / L) at 98 atom-% ¹⁵N was injected into each PVC core. This was performed during period of high precipitation (Figure 1) when water tables in the PVB were near the surface of the soil. Injections were made using 40 cm side-port needles, evenly distributing solution between 30 cm depth and the soil surface (Hart et al., 1994). Twenty injections were made per core, and were evenly distributed over the surface area of the cores to the extent possible. Points of injection were marked at the soil surface. Solution injections were equivalent to 11.1 kg N per ha and 20.4 mm of rainfall, and thus simulated realistic hydrologic NO₃ and water inputs. Soil solution NO₃-N concentrations ranged from 10-30 ppm, within the range of soil solution concentrations measured upslope of these PVB and presumably in subsurface flow entering these PVB (Zhou et al., 2010).

Soil samples (2 cm diameter to 30 cm depth) were collected on the day of isotope addition to determine initial soil NO₃ concentrations in the PVC cores. Samples were taken from outside of vegetated cores but inside of de-vegetated cores before isotope addition. Subsamples were

extracted in 2M KCl (5:1 extract: soil ratio). Extracts were filtered and frozen until analysis. Soil $\text{NO}_3\text{-N} + \text{NO}_2\text{-N}$ (hereafter $\text{NO}_3\text{-N}$) concentrations were determined by colorimetric analysis using the Griess-Ilosvay reaction with VCl_3 as a reducing agent (Hood-Nowotny et al., 2010). Soil gravimetric water content (VWC) to 30 cm depth was determined by drying subsamples of fresh soil at 105 °C. Bulk density (BD) for the watershed soils were taken from Gutierrez-Lopez (2012). Extract $\text{NO}_3\text{-N}$ concentrations and BD values were used to calculate the total mass of $\text{NO}_3\text{-N}$ in each core, and VWC data were converted to WFPS using these BD values.

Soil Collection

To measure recovery of ^{15}N in SOM pools 7 d after addition, soil cores (2 cm diameter to 30 cm depth) were collected from points of isotope injection within the PVC cores on 22 June 2011. To measure recovery of ^{15}N in SON pools over the full growing season, the PVC cores and all soil inside were collected on 29 (Watershed 1) and 31 Oct 2011 (Watersheds 2 and 3), which corresponded to 136 and 138 d after isotope addition (hereafter 137 d). After sampling on both days, soil was stored at 4 °C until processing.

Salt-extractable N and Microbial Biomass

Within 1 week of each soil collection, a sequential extraction procedure adapted from Holmes et al. (2003) was performed on 40 g subsamples of fresh soil to remove extractable C and N and microbial biomass. Soils were sieved to 2 mm, extracted in 0.5M K_2SO_4 (5:1

extract: soil ratio), and left to settle overnight at 4 °C. Extracts were filtered over pre-leached Whatman 42 filter papers. Extract total N concentrations were measured with a TOC-L CPN (Shimadzu, Kyoto, Japan) to determine and salt-extractable N, which includes NO₃-N, NH₄-N, and salt-extractable organic N (SEON). Extract NO₃-N concentrations were determined as described above, and extract NH₄-N concentrations were determined with the Berthelot reaction (Hood-Nowotny et al., 2010). Enrichment of salt-extractable N with ¹⁵N was not determined.

To extract microbial biomass, extracted soils were fumigated with chloroform (CHCl₃) using a direct fumigation procedure adapted from Witt et al. (2000) and Perakis and Hedin (2001). A 4.5 mL aliquot of CHCl₃ was pipetted directly onto each soil in the high-density polyethylene (HDPE) bottles. Bottles were immediately agitated by hand to spread the soil slurry across the bottle surface and maximize penetration of CHCl₃ into the soil pores. Fumigated soils and blanks were kept at room temperature in the dark for 5 d. After fumigation was complete, soils were subjected to vacuum venting to vaporize CHCl₃. Soils were then extracted in 0.5M K₂SO₄ as described above. Enrichment of microbial biomass N with ¹⁵N was not determined. Extracted soils were left to air-dry under the fume hood; once dry, they were stored at room temperature until further analysis.

Soil Organic Matter Analyses

Retention of ¹⁵N in POM and MAOM at 7 d and 137 d was measured in soils that had been sequentially extracted and air-dried. To separate POM and MAOM, subsamples were placed

in 0.08 M Na hexametaphosphate solution (4:1 solution: soil ratio) and shaken on a reciprocal shaker for ~20 hrs. Soil slurries were then emptied onto a 0.53 μm sieve and rinsed until all silt + clay + MAOM had washed through, with only sand + POM remaining on the sieve. Both fractions were dried at 65 °C. Subsamples (20-60 mg) of dried soil fractions were sent for analysis at the Stable Isotope Facility at University of California, Davis, CA, USA. Total C and N content and ^{15}N enrichment of the soil fractions were determined with an elemental analyzer (Elementar Analysensysteme GmbH, Hanau, Germany) interfaced to a PDZ Europa 20-20 isotope ratio mass spectrometer (Sercon Ltd., Cheshire, UK) at the Stable Isotope Facility at University of California, Davis, CA, USA.

Plant Biomass Collection and Analysis

Plant aboveground biomass (AGBM) from within the vegetated cores was harvested 137 d after isotope addition. Aboveground biomass was immediately set at 35 °C, subsequently oven-dried at 60 °C for 4 days, and pulverized. Rhizomes, root mats, and roots >2 mm in diameter that would not be represented by random sampling (hereafter “course roots”) were collected from air-dried soil from the cores collected at 137 d. Course roots were washed in 0.01 M CaCl_2 three times for 5 minutes to remove mineral N from root surfaces, oven-dried at 60 °C for 48 hrs, and weighed. Subsamples (~800 g) of air-dried bulk soil were collected for fine root analyses. Soils were placed in mesh containers (0.28 mm pore size) and washed in an elutriator to remove silt, clay, and fine sand. Fine roots and non-root litter were floated out of the remaining course sand, and non-root litter was removed manually. Fine roots were re-washed to remove remaining soil particles, dried at 60 °C for 48 hrs, and then pulverized.

Subsamples (3-5 mg) of pulverized AGBM, course root biomass (CRBM), and fine root biomass (FRBM) were sent for determination of total C and N and enrichment by ^{15}N at the Stable Isotope Facility at the University of California, Davis, as described above.

Denitrification Enzyme Assay

Soil samples (2 cm diameter to 30 cm depth) for denitrification enzyme assay (DEA) were collected on 25 June 2011. Soils were collected from the proximity of the PVC cores and from the equivalent topographic position in adjacent watersheds managed entirely as row crops (soybean in 2011). Soil $\text{NO}_3\text{-N}$ concentrations at time of sampling were determined by extracting in 2M KCl, as described above. To determine soil dissolved organic C (DOC) concentrations, fresh soil subsamples were extracted in 0.01 M CaCl_2 (2:1 extract: soil ratio), shaken orbitally at 160 rpm for 12 min, centrifuged at 2200 rpm for 15 min, and filtered to 0.45 μm . Extract C concentrations were measured by a Shimadzu TOC-L CPN.

Denitrification enzyme assay (DEA) was performed on fresh soil subsamples. The DEA protocol was based on Tiedje (1994). Soils were placed in 310 mL glass bottles and saturated in solution of 1 mM glucose ($\text{C}_6\text{H}_{12}\text{O}_6$) and KNO_3 . Soil slurries were flushed with He, evacuated repeatedly, and received 30 mL of acetylene (C_2H_2) gas. Soil slurries were then shaken to maintain gas equilibrium with the bottle headspaces. Gas samples were taken from the bottle headspaces at 0, 30, 60 and 90 min. Nitrous oxide (N_2O) concentrations in gas samples were determined with a gas chromatograph with an electron capture detector at

325°C. Gas species separation was accomplished with stainless steel columns packed with Haysep D and maintained at 50°C, using N₂ as carrier gas. Flux rates of N₂O from soil slurries were considered to represent denitrification enzyme potential. Only linear portions of fluxes were included in rate calculations (Tiedje, 1994).

Statistical Analyses

Statistical analyses were performed using PROC GLM in SAS (SAS Institute Inc., Cary, NC, USA). Watersheds were considered blocks; time of collection and vegetation treatment (for the PVC core soil data) and vegetation type (perennial vs. row crop) for the DEA data were considered fixed categorical factors.

Results

Soil Mineral N and Organic Matter

Initial soil NO₃-N concentrations in de-vegetated cores were lower than in vegetated cores ($p = 0.007$) and more variable (Table 1). Soil water contents were high (0.70-1.0 cm³ cm⁻³ WFPS to 30 cm depth) on the day of and 7 d after isotope addition, and did not differ between vegetation treatments. Soil NH₄-N and NO₃-N concentrations were greater in de-vegetated than in vegetated cores at both sampling times (Table 2).

Soil POM-N concentration increased between 7 d and 137 d, but there was an interactive effect of vegetation removal with time (Figure 2). Soil POM-N was greater in de-vegetated than vegetated cores at 7 d; however, POM-N increased from 7 d to 137 d in vegetated cores, but not in de-vegetated cores. Carbon: nitrogen ratio of POM decreased between 7 d and 137 d, but did not differ between vegetation treatments (Figure 2). Soil MAOM-N and MAOM-C concentrations increased from 7 d to 137 d across vegetation treatments, but there was no effect of vegetation treatment (Figure 3). Across vegetation treatments, MAOM C:N ratio increased from 7 d (mean=10.26) to 137 d (mean=10.40; $p = 0.055$).

Greater percentages of tracer $^{15}\text{NO}_3\text{-N}$ were recovered as MAOM-N than POM-N both 7 and 137 d after $^{15}\text{NO}_3$ addition (Figure 4). Percentage of $^{15}\text{NO}_3\text{-N}$ recovered as POM-N and MAOM-N did not differ between vegetation treatments at 7 d. However, at 137 d, greater percentages of $^{15}\text{NO}_3\text{-N}$ were recovered as POM-N and MAOM-N in vegetated than in de-vegetated cores (though this interaction was not significant for MAOM-N) (Figure 4). Percent of ^{15}N tracer recovered in POM + MAOM ranged from 3.6 to 10.1% (Figure 5).

Plant Biomass and Unrecovered ^{15}N

Tracer ^{15}N was recovered in AGBM, CRBM, and FRBM in all vegetated cores (Figure 5). Total % of ^{15}N tracer in plant biomass ranged from 4.2 to 19.5% (Figure 5). Percent ^{15}N recovered in AGBM was more variable than percent recovered in roots, and correlated positively with AGBM-N and negatively with AGBM C:N (data not shown). Similar trends were not found for percent ^{15}N recovered in roots. Total percent of tracer ^{15}N unrecovered in

plant biomass and SOM pools ranged from 70.4 to 92% (Figure 5). Percent of tracer ^{15}N unrecovered correlated negatively with % recovered in AGBM ($R^2 = 0.84$; data not shown).

Denitrification Enzyme Assay

Soil $\text{NO}_3\text{-N}$ concentrations were greater in row crop than PVB soils from equivalent landscape positions (Figure 6). However, denitrification enzyme potential showed the opposite trend, with DEA >50% higher in PVB compared to row crop soils, though the difference was not statistically significant (Figure 6). Water-filled pore space and DOC concentrations were greater in prairie than soybean soils (Figure 6).

Discussion

Retention of N in SOM

Previous data had shown contrasting soil C and N dynamics in watersheds with and without PVB. At the toeslope landscape position, organic C and N had increased and C:N had decreased in PVB soils since implementation, while organic N had decreased and C had not changed in row crop soils since conversion from brome (Pérez-Suárez, *unpublished data*). These data indicated that organic N had been accumulating in PVB soils but not row crop soils since treatment implementation, suggesting that SON could be a potentially significant sink for N inputs to these watersheds.

In this study, both POM-N and MAOM-N increased between mid-June and late October in cores with growing vegetation, indicating that these soils had accumulated organic N.

However, in de-vegetated cores, only MAOM-N increased over this time period.

Incorporation of $^{15}\text{NO}_3\text{-N}$ into POM and MAOM occurred rapidly (within 7 d of input) and independently of vegetation treatment. Similarly, Matheson et al. (2002) observed $^{15}\text{NO}_3\text{-N}$ incorporation into SOM within 24 d not to differ between planted and unplanted PVB soils.

However, in these soils, vegetation did affect longer-term recovery of $^{15}\text{NO}_3\text{-N}$ in SOM pools. In de-vegetated cores, amount of tracer ^{15}N recovered in POM and MAOM did not differ between 7 d and 137 d after addition. In contrast, the presence of growing perennial vegetation increased recovery of ^{15}N in both SOM pools at 137 d compared to 7 d, though this result was only statistically significant for POM. These results indicate that, while incorporation of $\text{NO}_3\text{-N}$ into both POM and MAOM occurred in these soils independently of plant growth, plant growth increased retention of $\text{NO}_3\text{-N}$ in both pools over the course of the growing season.

However, while these results show that $\text{NO}_3\text{-N}$ inputs were retained in SOM, and this retention was enhanced by plant growth, SOM was nevertheless a relatively minor sink for $\text{NO}_3\text{-N}$ inputs. The amount of ^{15}N recovered in SOM pools of vegetated cores after 137 d represented <10% of $^{15}\text{NO}_3\text{-N}$ inputs (Figure 5). Matheson et al. (2002) also found SOM to be a minor sink for $\text{NO}_3\text{-N}$ in PVB soils.

Retention of N in Plant Biomass

In this study, AGBM at the end of the growing season was the most variable sink for $^{15}\text{NO}_3\text{-N}$ inputs, while relatively consistent amounts were recovered in roots. Because the amount of ^{15}N retained in SOM and roots were relatively consistent, recovery in AGBM correlated closely and negatively with the amount of ^{15}N that was unrecovered. However, similarly to SOM, plant biomass at 137 d represented a minor sink for $\text{NO}_3\text{-N}$ inputs; total ^{15}N recovered in plant biomass represented <20% of $^{15}\text{NO}_3\text{-N}$ inputs, similar to results found by Matheson et al. (2002). Since only 3-7% of ^{15}N was recovered in AGBM at 137 d (Figure 5), late-fall harvest of AGBM from these watersheds is not likely to provide a major sink for $\text{NO}_3\text{-N}$ inputs. Total uptake of ^{15}N by vegetation was not determined in this study, and may represent a greater percentage of ^{15}N inputs. Plant biomass represents a temporary pool for N inputs, and N retained in herbaceous plant biomass would be expected to either mineralize or become SON on the scale of years to decades (Hefting et al., 2005). Further studies are needed to determine the long-term fate of N retained in plant biomass pools.

Unrecovered N and Denitrification as an N Sink

The majority of $^{15}\text{NO}_3\text{-N}$ inputs to these soils were not recovered in SOM or plant biomass. The amount of $^{15}\text{NO}_3$ leached out of the PVC cores (to depths >30 cm) could not be determined in this study. However, since subsurface NO_3 losses from watersheds with PVB have been substantially decreased compared to watersheds without PVB (Zhou et al., 2010), subsurface leaching appears not to be a major loss pathway for NO_3 in these watersheds. Therefore, even if substantial amounts of $^{15}\text{NO}_3$ had been lost from the soils within the PVC

cores by leaching, whole-watershed data indicate that the majority of this $^{15}\text{NO}_3$ would not have been lost from the PVB in subsurface flow.

The lack of recovery of ^{15}N in SOM and plant biomass and demonstrated decrease in subsurface dissolved NO_3 losses from these watersheds together indicate that denitrification is the major sink for NO_3 inputs to these PVB. Denitrification has been found to be the major NO_3 sink in many riparian PVB (e.g. Martin et al., 1999; Matheson et al., 2002). If denitrification had accounted for 100% of unrecovered $^{15}\text{NO}_3\text{-N}$ inputs in this study, it would be equivalent to 7.3-9.6 kg $\text{NO}_3\text{-N}$ denitrified ha^{-1} , which is within observed denitrification rates in herbaceous PVB during summer months (Hefting et al., 2003, 2004) as well as other PVB systems (e.g. Hanson et al., 1994b).

Recovery of ^{15}N in salt-extractable N and microbial biomass were not determined.

Concentrations of extractable N measured in the cores at 7 indicate that this pool could have accounted for as much as 30-100% of the unrecovered $^{15}\text{NO}_3\text{-N}$ at this time (data not shown). However, mineral and extractable organic N forms represent a very short-term sink for N inputs, and any ^{15}N recovered in these pools would be expected to enter another sink or be lost from the system on a short time scale. Microbial biomass has been shown to not function as a major sink for N inputs in PVB (Hanson et al., 1994a).

The results from the DEA showed that, despite substantially lower NO_3 concentrations, greater mean denitrification enzyme potentials were observed in PVB soils than row crop soils in each watershed, indicating that greater percentages of soil NO_3 are denitrified in PVB

than row crop soils. Greater denitrification enzyme potential corresponded to greater DOC concentrations, as well as slightly but consistently greater WFPS in PVB than row crop soils. Given that WFPS were high ($>0.70 \text{ cm}^3 \text{ cm}^{-3}$) in both vegetation types at the time of soil collection, greater availability of DOC substrate for denitrification more likely accounted for the difference in denitrification enzyme potential. Data from the DEA procedure cannot be used to quantify denitrification rates in the field, and, in this study, were only collected during one part of the season. Thus, further research is needed to verify the role of denitrification in decreasing NO_3 loss in subsurface flow from these watersheds. Additionally, since denitrification can produce nitrous oxide (N_2O), denitrification in PVB can potentially change an aquatic pollutant to an air pollutant (Verhoeven et al., 2006; Stevens and Quinton, 2009). Further research is needed to determine the composition of gaseous N ($\text{N}_2\text{O} / \text{N}_2$) produced by denitrification in these PVB.

Conclusion

These results indicate that denitrification is the major sink for $\text{NO}_3\text{-N}$ in PVB. While $\text{NO}_3\text{-N}$ inputs are retained in SOM and plant biomass, the magnitudes of these sinks are small. Since denitrification can remove NO_3 from the soil indefinitely under favorable conditions, these results indicate that initial reductions in subsurface NO_3 losses observed in these watersheds will likely continue in the long term. Future research is needed to determine the nature of gaseous N products and whether or not similar results would be observed in comparable sites with varying hydrological and topographic features.

Acknowledgements

The authors thank the following individuals for their assistance: Pauline Drobney and the staff of Neal Smith NWR for their support of this project; Carl Pedersen for installing the PVC cores; Chris Witte for support with field work and equipment throughout the project; David Sundberg for providing use of the elutriator for root analysis; Teresita Chua for providing use of a microbalance; Danielle Wilson for providing use of strainers for roots; Xiaobo Zhou for preliminary and supporting data from the site; and Kirsten Hofmockel for insight and advice in the implementation and interpretation of this research. Funding for the research site was provided by the Leopold Center for Sustainable Agriculture, Iowa State University College of Agriculture and Life Sciences, USDA Forest Service Northern Research Station, and the Iowa Department of Agriculture and Land Stewardship.

References

- Aber JD, Nadelhoffer KJ, Steudler P, Melillo JM. 1989. Nitrogen saturation in northern forest ecosystems. *BioScience* 39: 378-386.
- Aber J, McDowell W, Nadelhoffer K, et al. 1998. Nitrogen saturation in temperate forest ecosystems: hypotheses revisited. *BioScience* 48: 921- 934.
- Alexander RB, Smith RA, Schwarz GE, Boyer EW, Nolan JV, Brakebill JW. 2008. Differences in phosphorus and nitrogen delivery to the Gulf of Mexico from the Mississippi River basin. *Environmental Science and Technology* 42: 822-830.
- Baker MA, Vervier P. 2004. Hydrological variability, organic matter supply and denitrification in the Garonne River ecosystem. *Freshwater Biology* 49: 181-190.
- Barrett JE, Burke JC. 2002. Nitrogen retention in semiarid ecosystems across a soil organic-matter gradient. *Ecological Applications* 12: 878-890.

- Bedard-Haughn A, Tate KW, van Kessel C. 2004. Using nitrogen-15 to quantify vegetative buffer effectiveness for sequestering nitrogen in runoff. *Journal of Environmental Quality* 33: 2252-2262.
- Booth MS, Stark JM, Rastetter E. 2005. Controls on nitrogen cycling in terrestrial ecosystems: a synthetic analysis of literature data. *Ecological Monographs* 75: 139-157.
- Carpenter SR, Caraco NF, Correll DL, Howarth RW, Sharpley AN, Smith VH. 1998. Nonpoint pollution of surface waters with phosphorus and nitrogen. *Ecological Applications* 8: 559-568.
- Castellano MJ, Kaye JP, Lin H, Schmidt JP. 2012. Linking carbon saturation concepts to nitrogen saturation and retention. *Ecosystems* 15: 175-187.
- David MB, Drinkwater LE, McIsaac GF. 2010. Sources of nitrate yields in the Mississippi River basin. *Journal of Environmental Quality* 39: 1657-1667.
- Dell CJ, Williams MA, Rice CW. 2005. Partitioning of nitrogen over five growing seasons in tallgrass prairie. *Ecology* 86: 1280-1287.
- Feng W, Plante AF, Six J. 2012. Improving estimates of maximal organic carbon stabilization by fine soil particles. *Biogeochemistry* 95: 1-13.
- Gutierrez-Lopez J. 2012. Soil moisture dynamics in agriculturally-dominated landscapes after the introduction of native prairie vegetation. *Graduate Theses and Dissertations, Iowa State University*. Paper 12337.
- Hanson GC, Groffman PM, Gold AT. 1994a. Symptoms of nitrogen saturation in a riparian wetland. *Ecological Applications* 4: 750-756.
- Hanson GC, Groffman PM, Gold AT. 1994b. Denitrification in riparian wetlands receiving high and low groundwater nitrate inputs. *Journal of Environmental Quality* 23: 917-922.
- Hart SC, Stark JM, Davidson EA, Firestone MK. 1994. Nitrogen mineralization, immobilization, and nitrification. In: Weaver, R.W., Angle, J.S., Bottomley, P.S.

- (Eds.), *Methods of Soil Analysis, Part 2 – Microbiological and Biochemical Properties*. Soil Science Society of America, Madison, WI, USA, pp. 985-1018.
- Hassink J. 1997. The capacity of soils to preserve organic C and N by their association with clay and silt particles. *Plant and Soil* 191: 77-87.
- Haycock NE, Pinay G. 1993. Groundwater nitrate dynamics in grass and poplar vegetated riparian buffer strips during the winter. *Journal of Environmental Quality* 22: 273-278.
- Hefting MM, Bobbink R, de Caluwe H. 2003. Nitrous oxide emission and denitrification in chronically nitrate-loaded riparian buffer zones. *Journal of Environmental Quality* 32: 1194-1203.
- Hefting MM, Clement JC, Dowrick D, Cosandey AC, Bernal S, Cimpian C, Tatur A, Burt TP, Pinay G. 2004. Water table elevation controls on soil nitrogen cycling in riparian wetlands along a European climatic gradient. *Biogeochemistry* 67: 113-134.
- Hefting MM, Clement J-C, Bienkowski P, Dowrick D, Guenat C, Butturini A, Topa S, Pinay G, Verhoeven JTA. 2005. The role of vegetation and litter in the nitrogen dynamics of riparian buffer zones in Europe. *Ecological Engineering* 24: 465-482.
- Hill AR. 1996. Nitrate removal in stream riparian zones. *Journal of Environmental Quality* 25: 743-755.
- Hirsh SM. 2012. Diversifying monoculture crops by incorporating prairie buffer strips. Graduate Theses and Dissertations, Iowa State University. Paper 12343.
- Holmes WE, Zak DR, Pregitzer DS, King JS. 2003. Soil nitrogen transformations under *Populus tremuloides*, *Betula papyrifera* and *Acer saccharum* following 3 years exposure to elevated CO₂ and O₃. *Global Change Biology* 9: 1743-1750.
- Hood-Nowotny R, Umana NH-N, Inselbacher E, Oswald-Lachouani P, Wanek W. 2010. Alternative methods for measuring inorganic, organic, and total dissolved nitrogen in soil. *Soil Science Society of America Journal* 74: 1018-1027.
- Kaye J, Barrett J, Burke IC. 2002. Stable nitrogen and carbon pools in grassland soils of variable texture and carbon content. *Ecosystems* 5: 461-471.
- Kleber M, Sollins P, Sutton R. 2007. A conceptual model of organo-mineral interactions in

- soils: self-assembly of organic molecular fragments into zonal structures on mineral surfaces. *Biogeochemistry* 85: 9-24.
- Lowrance R. 1992. Groundwater nitrate and denitrification in a coastal plain riparian forest. *Journal of Environmental Quality* 21: 401-405.
- Martin TL, Kaushik NK, Trevors JT, Whiteley HR. 1999. Review: denitrification in temperate climate riparian zones. *Water, Air and Soil Pollution* 111: 171-186.
- Matheson FE, Nguyen ML, Cooper AB, Burt TP, Bull DC. 2002. Fate of ¹⁵N-nitrate in unplanted, planted and harvested riparian wetland soil microcosms. *Ecological Engineering* 19: 249-264.
- Mayer PM, Reynolds SK, McCutchen MD, Canfield TJ. 2007. Meta-analysis of nitrogen removal in riparian buffers. *Journal of Environmental Quality* 36: 1172-1180.
- Perakis SS, Hedin LO. 2001. Fluxes and fates of nitrogen in soil of an unpolluted old-growth temperate forest, southern Chile. *Ecology* 82: 2245-2260.
- Schade JD, Fisher SG, Grimm NB, Seddon JA. 2001. The influence of a riparian shrub on nitrogen cycling in a Sonoran desert stream. *Ecology* 82: 3363-3376.
- Schade JD, Lewis DB. 2006. Plasticity in resource allocation and nitrogen-use efficiency in riparian vegetation: implications for nitrogen retention. *Ecosystems* 9: 740-755.
- Six J, Conant RT, Paul EA, Paustian K. 2002. Stabilization mechanisms of soil organic matter: implications for C-saturation of soils. *Plant and Soil* 241: 155-176.
- Stevens CL, Quinton JN. 2009. Diffuse pollution swapping in arable agricultural systems. *Critical Reviews in Environmental Science and Technology* 36: 478-520.
- Stewart CE, Paustian K, Conant RT, Plante AF, Six J. 2007. Soil carbon saturation: concept, evidence and evaluation. *Biogeochemistry* 86: 19-31.
- Townsend AR, Howarth RW, Bazzaz FA, Booth MS, Cleveland CC, Collinge SK, Dobson AP, Epstein PR, Holland EA, Keeney DR, Mallin MA, Rogers CA, Wayne P, Wolfe AH. 2003. Human health effects of a changing global nitrogen cycle. *Frontiers in Ecology and Environment* 1: 240-246.

- Verchot LV, Franklin EC, Gilliam JW. 1997. Nitrogen cycling in Piedmont vegetated filter zones: II. Subsurface nitrate removal. *Journal of Environmental Quality* 26: 337-347.
- Verhoeven JTA, Arheimer B, Yin C, Hefting MM. 1996. Regional and global concerns over wetlands and water quality. *Trends in Ecology and Evolution* 21: 96-103.
- Vitousek PM, Reiners WA. 1975. Ecosystem succession and nutrient retention: a hypothesis. *BioScience* 25: 376-381.
- Vought LB-M, Dahl J, Pedersen CL, Lacoursière JO. 1994. Nutrient retention in riparian ecotones. *Ambio* 23: 342-348.
- Witt C, Gaunt JL, Galicia CC, Ottow JCG, Neue H-U. 2000. A rapid chloroform-fumigation extraction method for measuring soil microbial biomass carbon and nitrogen in flooded rice soils. *Biology and Fertility of Soils* 30: 510-519.
- Zhou X, Helmers MJ, Asbjornsen H, Kolka R, Tomer MD. 2010. Perennial filter strips reduce nitrate levels in soil and shallow groundwater after grassland-to-cropland conversion. *Journal of Environmental Quality* 39: 2006-2015.

Table 2.1. Soil NO₃-N concentrations by watershed and vegetation treatment immediately before and immediately following isotope addition. Values in parentheses are standard deviations of 3 replicates.

Watershed	Vegetation Treatment	mg NO ₃ -N kg ⁻¹ soil	
		(before)	(after)
1	Vegetated	0.69 (0.32)	3.45 (0.32)
	De-vegetated	2.83 (1.75)	4.58 (1.75)
2	Vegetated	0.45 (0.20)	3.33 (0.20)
	De-vegetated	1.94 (2.57)	4.82 (2.57)
3	Vegetated	0.38 (0.25)	3.25 (0.25)
	De-vegetated	3.99 (4.26)	6.86 (4.26)

Table 2.2. Soil NH₄-N and NO₃-N concentrations by watershed and vegetation treatment 7 and 137 days after isotope addition. Values in parentheses are standard deviations of 3 replicates.

Watershed	Vegetation Treatment	mg NH ₄ -N kg ⁻¹ soil		mg NO ₃ -N kg ⁻¹ soil	
		7 d	137 d	7 d	137 d
1	Vegetated	1.10 (0.09)	2.27 (0.56)	0.56 (0.40)	0.49 (0.51)
	De-vegetated	1.89 (0.99)	2.99 (0.56)	1.83 (1.70)	4.10 (1.47)
2	Vegetated	0.76 (0.65)	1.34 (0.99)	0.61 (0.40)	2.68 (3.03)
	De-vegetated	2.31 (1.72)	3.95 (1.78)	1.31 (0.62)	5.22 (1.59)
3	Vegetated	0.78 (0.68)	5.45 (0.23)	0.51 (0.25)	0.97 (0.80)
	De-vegetated	2.43 (1.46)	5.55 (1.61)	2.27 (1.74)	2.33 (2.33)




Figure 2.1. Precipitation at the research site in 2011, showing dates for ^{15}N application (June 15), collection of soils for DEA (June 25), and final collection of soils and biomass in the cores (October 29 and 31).




Figure 2.2. Soil particulate organic matter nitrogen (POM-N) concentrations and carbon: nitrogen ratio (POM C:N) by vegetation treatment and time after isotope addition (7 and 137 d), with analysis of variance results. Error bars show standard error of means of 9 replicates.




Figure 2.3. Soil mineral-associated organic matter nitrogen (MAOM-N) and carbon (MAOM-C) concentrations by vegetation treatment and time after isotope addition (7 and 137 d), with analysis of variance results. Error bars show standard error of means of 9 replicates.




Figure 2.4. Percent recovery of ^{15}N tracer in particulate organic matter (POM) and mineral-associated organic matter (MAOM) by vegetation treatment and time after isotope addition (7 and 137 d), with analysis of variance results shown for POM and MAOM. Error bars show standard errors of means of 9 replicates.




Figure 2.5. Percent of ^{15}N tracer recovered in aboveground biomass (AGBM), course root biomass (CRBM), fine root biomass (FRBM), particulate organic matter (POM), mineral-associated organic matter (MAOM), and unrecovered in vegetated cores 137 days after ^{15}N addition. Note break and change of scale in the y-axis. Error bars show standard errors of means of 9 replicates.




Figure 2.6. Comparison of A) NO₃-N concentrations, B) denitrification enzyme potential, C) water-filled pore space (WFPS), and D) dissolved organic carbon (DOC) concentrations in soils from equivalent topographic positions in perennial vegetation buffers and row crops (soybean) collected 25 June 2011. Analysis of variance p-values are for comparisons between vegetation types. Error bars show standard errors of means of 9 replicates.

CHAPTER 3. COVER CROP EFFECTS ON NITROUS OXIDE EMISSIONS FROM A MAIZE-BASED CROPPING SYSTEM: INTERACTION OF CARBON AND NITROGEN INPUTS

A paper submitted to the journal *Soil Biology and Biochemistry*

David C. Mitchell⁵, Michael J. Castellano⁶, John E. Sawyer⁷, Jose L. Pantoja⁸

Abstract

Nitrous oxide (N₂O) emissions from cultivated soils often increase with N fertilizer inputs and soil nitrate (NO₃) concentrations. The growth of cover crops from fall to spring can reduce soil NO₃ concentrations and has been promoted as a strategy to decrease N₂O emissions. However, mineralizable carbon (C) availability can be a more important control on N₂O emissions than NO₃ in inorganic N-rich soils, and cover crop residue provides a mineralizable C input. We measured the effect of a winter rye (*Secale cereale* L.) cover crop on soil N₂O emissions from a maize (*Zea mays* L.)-based cropping system treated with three N fertilizer rates (0, 135, and 225 kg N ha⁻¹) at a site in Iowa, USA. In addition, we conducted a laboratory incubation to determine if N₂O emissions from soils from these treatments were limited by mineralizable C or NO₃. The rye cover crop decreased soil NO₃ concentrations at all N rates. Although the cover crop decreased N₂O emissions at 0 kg fertilizer N ha⁻¹, it increased N₂O emissions from fertilized treatments. Results from

⁵ Graduate student, Department of Agronomy, ISU. Primary researcher and author of the paper.

⁶ Assistant professor, Department of Agronomy, ISU. Major professor of DC Mitchell.

⁷ Professor, Department of Agronomy, ISU. Contributed intellectually to this paper.

⁸ Graduate student, Department of Agronomy, ISU. Contributed intellectually to this paper.

laboratory incubations were consistent with field data; N₂O emissions from fertilized treatments did not increase with added NO₃, but did increase with added glucose. These results indicate that mineralizable C inputs can limit N₂O emissions and that C inputs from the rye cover crop likely increased N₂O emissions from fertilized soils. Mineralizable C availability should be considered in future evaluations of cover crop effects on N₂O emissions, particularly as cover crops are evaluated as a potential strategy to mitigate agricultural N₂O emissions.

Keywords: Nitrous oxide, cover crops, nitrate, mineralizable carbon, denitrification

Introduction

Atmospheric concentration of the potent greenhouse gas nitrous oxide (N₂O) has increased 40-50% since pre-industrial times as a consequence of human activity (Smith et al., 2007). Agricultural soils are the largest anthropogenic source of N₂O (Reay et al., 2012). The microbial processes nitrification and denitrification, as well as abiotic processes, produce N₂O in soils (Bremner, 1997). In cultivated soils, denitrification tends to be the dominant process producing N₂O (Ostrom et al., 2010). In denitrification, NO₃ is reduced to N₂O during organic carbon (C) oxidation in the absence of oxygen (O₂). Nitrous oxide can be further reduced to N₂ before diffusing from the soil; however, high NO₃ availability inhibits N₂O reduction (Blackmer and Bremner, 1978; Firestone and Davidson, 1989). Thus, emission of denitrification-derived N₂O is controlled by anaerobicity, the availability NO₃

and mineralizable C substrates, and the amount of N₂O reduced to N₂ before leaving the soil (Firestone and Davidson, 1989).

Laboratory studies of denitrification (Myrold and Tiedje, 1985; Lalisse-Grundmann et al., 1988) and field studies of N₂O emissions (Sehy et al., 2003; van Groenigen et al., 2004; Adviento-Borbe et al., 2007) have shown that these processes are not necessarily limited by NO₃ concentration in fertilized soils. In soils with high NO₃ concentrations and low O₂ availability, mineralizable C availability can control denitrification (Burford and Bremner, 1975; Bijay-Singh et al., 1988) and mineralizable C availability has been shown to limit N₂O emission from NO₃-rich (30 - 100 mg NO₃-N kg⁻¹ soil) cultivated soils (e.g. Weier et al., 1993; McKenney et al., 1995; Gillam et al., 2008), even with high soil organic C content (Sainz Rozas et al., 2001; Sánchez-Martín et al., 2008).

Overwintering non-legume cover crops can decrease NO₃ concentrations in agricultural soils through N uptake during growth and immobilization during residue decomposition (Thorup-Kristensen, 2003). By decreasing the soil NO₃ pool, these cover crops can also decrease soil N₂O emissions (Baggs et al., 2000). However, some studies have found that cover crops increase (Petersen et al., 2011) or have no consistent effect on N₂O emissions (e.g. Jarecki et al., 2009; Smith et al., 2011). When cover crops are killed shortly before N application, mineralizable C from their residue may stimulate denitrification and N₂O emissions (Sarkodie-Addo et al., 2003; Petersen et al., 2011).

Together, inconsistent effects of cover crops on N₂O emissions and evidence for C limitation of agricultural N₂O emissions indicate that cover crop effects on the availability of C as well as NO₃ can affect N₂O emissions. An improved understanding of the relationship between cover crops and N₂O emissions is particularly important as cover crops are increasingly promoted as a greenhouse gas mitigation strategy (Eagle and Olander, 2012). The goal of this study was to determine how cover crop C inputs interact with soil NO₃ to affect N₂O emissions in a maize-soybean [*Glycine max* (L.) Merr.] crop rotation in Iowa, USA. We used a coupled field and laboratory experimental approach. In the field, we measured the effects of three N fertilizer rates and a rye cover crop on soil N₂O emissions in the maize phase of the rotation. In the laboratory, we incubated soils from the field experiment with glucose and NO₃ additions to determine biochemical limitations on N₂O emissions as related to field observations.

Methods

Field Study

A field study was conducted at the Iowa State University Agricultural Engineering and Agronomy Ames Research Farm in Boone County, Iowa, USA (42.02N, 93.77W). Long-term mean annual temperature and precipitation at this location are 9.4 °C and 872 mm yr⁻¹. The soil at this site is Clarion loam series (fine-loamy, mixed, superactive, mesic Typic Hapludolls), with pH 6.4 in water (1:1 soil: water ratio), 2.4% total C, and 0.2% total N. The site was managed as a no-till maize-soybean rotation and treated in a split-plot design with

four replicates. Presence or absence of a winter rye cover crop was the main plot and N fertilizer rate for maize was the sub-plot. Of the six N rates at this site, those included in this study were 0, 135, and 225 kg N ha⁻¹ (hereafter N0, N135, and N225). No N was applied to soybean. Cover crop treatments were begun in fall 2008 and N rate treatments in spring 2009. Individual plot size was 6.1 m (8 maize rows) × 15.2 m length. The experimental site for this study was primarily established to better determine the economical optimum N rate (EONR) for maize following soybean with and without a rye cover crop. The long-term optimal N rate for this rotation in Iowa is 152 kg N ha⁻¹ at a fertilizer to grain price ratio of 0.1, according to the Corn Nitrogen Rate Calculator (Iowa State Univ. Agronomy Extension, 2012), a widely-used resource for N fertilizer recommendations in the USA (Sawyer et al., 2006; Robertson and Vitousek, 2009).

Rye was drill-seeded after soybean harvest on 5 Oct 2010 at 70 kg seed ha⁻¹ and killed with glyphosate [*N*-(phosphonomethyl)glycine] on 2 May 2011. Rye aboveground biomass was sampled before glyphosate application using a 0.09 m² frame at six random locations per replicate. Since soil NO₃ concentrations following 2010 soybean harvest did not differ between N rates, rye biomass was pooled from all N rates per replicate. Samples were dried at 60 °C, ground to pass a 2 mm sieve, and sub-sampled for total C analysis by dry combustion (LECO CHN-2000 analyzer, LECO Corp., St. Joseph, MI, USA). Maize was planted on 10 May 2011 at 76-cm row spacing with the planter equipped with residue cleaners and no-till coulters. The N fertilizer was urea ammonium nitrate (UAN, 32% N), side-dressed in bands to 15 cm depth between every other crop row on 19 May 2011. Nitrogen fertilizer was therefore highly concentrated in a small soil zone, as is typical for

injected N applied to maize. Maize grain was machine harvested on 5 Oct 2011 from the center four rows of each plot, with yield adjusted to 155 g kg^{-1} moisture.

Soil surface N_2O fluxes were measured approximately fortnightly from 11 Apr to 3 Oct 2011 using the static chamber method (Parkin and Venterea, 2010). Polyvinyl chloride (PVC) rings (25 cm diameter x 10 cm height) were placed in the plots to 5 cm depth. Two rings were installed in each plot in the following configuration: one ring directly over the N fertilizer band, the second ring overlapping the adjacent crop row and the next inter-row space (which received no N fertilizer). A vented PVC lid (25 cm diameter x 5 cm height) was used for a total chamber volume of 0.0049 m^3 . Change in N_2O concentration inside the chambers was analyzed *in situ* with a 1412 Infrared Photoacoustic Gas Monitoring System (Innova Air Tech Instruments, Ballerup, Denmark) (Iqbal et al., 2012). Fluxes were measured between 08:00 and 14:00 hrs to obtain flux rates proximate to the 24 hr period. Fluxes were measured in all 6 treatments in one replicate each day of measurement, with the treatment order randomized each day. During each measurement, N_2O concentration in the chamber was measured every 2 min over 14 min for a total of eight measurements including time zero. All eight measurements were used to calculate flux rates with a linear model. Flux rates with $R^2 < 0.65$ were considered zero. Flux rates from the two rings were weighted proportional to the area of the plot they represented. Cumulative N_2O emissions from the period of measurement were calculated by linear interpolation and numerical integration between sample times.

During each gas flux measurement, soil temperature at 5 cm depth was measured with a thermometer and volumetric water content (VWC) to 5 cm was measured with a TH300 theta probe (Dynamax Inc., Houston, TX, USA). Since soil bulk density did not vary with time or treatment over the period of measurement, VWC data were converted to water-filled pore space (WFPS) using the average bulk density (1.09 g cm^{-3}) for the site. One soil sample (2 cm diameter to 10 cm depth) was collected during each gas flux measurement corresponding to each gas flux ring location. A soil subsample was extracted in 2 M KCl (5:1 solution: soil ratio) by shaking for 1 hr at 180 rpm. Extracts were filtered through pre-leached Whatman 1 filter paper and frozen until analysis. Extract ($\text{NO}_3 + \text{NO}_2$) (hereafter NO_3) and NH_4 concentrations were measured in microplates using the Griess-Ilosvay reaction with VCl_3 as a reducing agent and the Berthelot reaction, respectively (Hood-Nowotny et al., 2010). Soil mineral N concentrations corresponding to the two gas flux rings were weighted proportional to the area of the plot they represented.

Incubation Study

Soil cores (2 cm diameter to 10 cm depth) were collected from the N fertilizer bands in all field treatments between 13 and 16 June 2011, during a period of high N_2O flux rates. Soils were air-dried ($\sim 25 \text{ }^\circ\text{C}$) upon returning to the lab and sieved to 4 mm. Initial NO_3 and NH_4 concentrations were determined in a subsample of air-dried soil using the methods described above.

Subsamples of 15 g air-dried soil from each field treatment were placed in 120 mL bottles. In a pilot study, soil water content of 90% water-holding capacity (WHC), or $\sim 0.80 \text{ cm}^3 \text{ cm}^{-3}$ WFPS, was determined to maximize N_2O emissions from these soils (data not shown). Soils were brought to this water content with one of 4 treatment solutions: de-ionized water, 1 mM KNO_3 in water, 1 mM glucose ($\text{C}_6\text{H}_{12}\text{O}_6$) in water, or 1 mM of both KNO_3 and glucose in water. Concentrations of KNO_3 and glucose in treatment solutions were based on the denitrification enzyme assay protocol (Tiedje 1994). Treatment solutions were applied factorially to soils from each field treatment from each replicate. Soils were then incubated in the dark at $\sim 22^\circ\text{C}$ with bottles open to laboratory air. Soils were maintained at 90% WHC by periodic re-application of the treatment solutions, though initial application accounted for 91% of water, NO_3 , and glucose added. Total additions over the 10 day incubation were 7.8 mg $\text{NO}_3\text{-N}$ and 40.3 mg glucose-C kg^{-1} soil.

Nitrous oxide emissions from the incubating soils were measured 1, 4, and 10 d after initial treatment solution application. Air samples were collected from the bottle headspaces at 0 and 30 min after sealing the bottles with rubber septa and stored in pre-evacuated Exetainer® vials (Labco Ltd., Lampeter, Ceredigion, UK). An equal volume of air was added to each bottle immediately after removing the 0 min samples to maintain headspace pressure.

Concentrations of N_2O in the vials were measured with a gas chromatograph (Agilent 7890, Santa Clara, CA, USA) operated with an electron capture detector at 350°C . Gas species separation was accomplished with stainless steel columns packed with Porapak Q, 80/100 mesh and maintained at 85°C . Carrier gas was 10% CH_4 and 90% Ar. Carbon dioxide production in the incubating soils was measured more frequently during the 10 day

incubation using a LI-7000 infrared gas analyzer (LI-COR, Omaha, NE, USA). Cumulative N₂O and CO₂ emissions were calculated by linear interpolation and numerical integration between sample times.

Statistical Analyses

All cumulative N₂O emission and soil NO₃ and NH₄ concentration data were log-transformed before analysis to meet the assumptions of normality and homogeneity of residuals. Soil temperature, moisture, and NO₃ and NH₄ concentration data from the field study were analyzed as repeated measures. All data were analyzed using PROC GLM in Statistical Analysis Software package (SAS Institute Inc., Cary, NC, USA).

Results

Field Study

Soil temperatures were highest in N0 but did not vary between the other N rates or cover crop treatments. Water-filled pore space did not differ between cover crop treatments or N rates. The rye cover crop had no effect on soil NH₄ concentrations (data not shown) but decreased NO₃ concentrations at all three N rates (Figure 1). Soil NO₃ concentrations were greater at N225 than N135 (Figure 1). Rye produced 160-345 kg C ha⁻¹ in aboveground biomass. Maize yields were greater at N225 than N135 in both cover crop treatments (Table 1).

Cover crop and N rate treatments interacted to affect N₂O emissions (Figure 2). The rye cover crop decreased cumulative N₂O emissions at N0, increased emissions at N135, and had no effect at N225. Without the cover crop, N₂O emissions were greater at N225 than N135. However, with the cover crop, emissions did not differ between these N rates. The fertilizer bands were the dominant source of N₂O from the fertilized treatments at this site despite the small portion of the plot area they covered. On average, fertilizer bands accounted for 18, 65, and 70% of total plot N₂O emissions at N0, N135, and N225, respectively.

Incubation

Initial NO₃ concentrations were lower in soils with the cover crop than without cover crop across all N fertilizer rates (Figure 3). Initial NO₃ in incubated soils were greater in soils from N225 than N135 and lowest in soils from N0 (Figure 3). Across NO₃ addition and glucose addition treatments, cumulative CO₂ emission from incubating soils did not differ between N fertilizer rate and cover crop treatments. Average CO₂ emissions from incubating soils over 10 days were 222.3 mg CO₂-C kg⁻¹ soil from soils with glucose added and 195.0 mg CO₂-C kg⁻¹ soil from soils without glucose added. In contrast, N fertilizer rate (field treatment) had a main effect on cumulative N₂O emissions from the incubated soils (Table 2). Cover crop and NO₃ addition treatments did not have main effects on N₂O emissions from incubating soils; however, there was an interaction between N fertilizer rate and NO₃ addition (Table 2). Across cover crop and glucose addition treatments, NO₃ addition increased N₂O emission from N0 soils but not N135 and N225 soils (Figure 4). In contrast,

there was a main effect of glucose addition on N₂O emission, as well as an interaction between cover crop, N fertilizer rate, and glucose addition treatments (Table 2, Figure 5). In soils from N0, glucose addition increased cumulative N₂O emission only in soils without cover crop. In soils from N135, glucose addition increased N₂O emissions in soils from both cover crop treatments. In soils from N225, glucose addition increased N₂O emissions in soils with cover crop but not without cover crop (Figure 5).

Discussion

Field N₂O Emissions

Nitrogen fertilizer is a major control on N₂O emissions from cultivated soils (Stehfest and Bouwman, 2006). Emissions of N₂O have been found to increase with N fertilizer rate both linearly (e.g. Mosier et al., 2006; Smith et al., 2011) and non-linearly (e.g. McSwiney and Robertson, 2005; Stehfest and Bouwman, 2006). However, other studies have not found consistent increases in N₂O emissions with N fertilizer rate (e.g. Sehy et al., 2003; van Groenigen et al., 2004; Adviento-Borbe et al., 2007). In this study, N₂O emissions were greater in fertilized treatments than N0, but the relationship between N₂O emission and N rate was not consistent between cover crop treatments. N₂O emissions were greater at N225 than N135 without the rye cover crop, but did not differ with the rye cover crop, despite greater soil NO₃ concentrations in N225 than N135 across cover crop treatments (Figure 1 and 2).

Since non-legume cover crops can decrease soil NO_3 concentrations through plant uptake and microbial immobilization during residue decomposition, they have potential to decrease N_2O emissions (Baggs et al., 2000; McSwiney et al., 2010). However, the rye cover crop in this study only decreased N_2O emissions at N_0 , despite decreasing soil NO_3 concentrations at all N rates (Figure 1 and 2). In contrast, the cover crop in this study increased N_2O emissions at N_{135} . Although the cover crop had no consistent effect on N_2O emissions at N_{225} , variability at N_{225} with cover crop was exceptionally high (Figure 2). These results indicate that N_2O emissions were not limited by soil NO_3 concentration in the fertilized treatments, even with a decreased NO_3 pool with the cover crop. Since soil moisture and temperature did not differ between cover crop treatments, the increase in N_2O emission with the rye cover crop at N_{135} cannot be explained by cover crop effects on soil temperature, moisture, or mineral N concentrations.

Limitation of N_2O emission by available C may partially explain the inconsistent relationships between N_2O emissions, N fertilizer rate, and mineral N concentrations. Cover crop residue inputs to NO_3 -rich soils provide mineralizable C substrate that can stimulate N_2O production (McKenney et al., 1995), especially when the cover crop kill coincides with N fertilizer application (Sarkodie-Addo et al., 2003; Petersen et al., 2011). In this study, the rye cover crop provided 160-345 kg C ha^{-1} in aboveground residue, in addition to root residue (not quantified in this study), 17 days before N application. A portion of this C was likely still available for microbial mineralization following N application. Thus, C inputs from cover crop residue may have influenced the increased N_2O emissions with the cover crop at N_{135} .

Incubation Treatments and Limitation on N₂O Emission

In the incubation, N₂O emission was limited by NO₃ availability in soils from N0 and by mineralizable C availability in soils from N135 and N225. This result agrees with previous studies that have demonstrated that denitrification and N₂O emission are limited by NO₃ availability in low-NO₃ soils and by mineralizable C availability in high-NO₃ soils (Weier et al., 1993; Gillam et al., 2008). Although initial NO₃ concentrations were lower in fertilized soils with the rye cover crop than without, NO₃ addition to these soils during incubation did not increase N₂O emissions (Table 2, Figure 3 and 4). In contrast, glucose addition did increase N₂O emissions from fertilized soils (Table 2, Figure 5), demonstrating that mineralizable C inputs, such as those from cover crop residue, can stimulate N₂O emissions. While C mineralization in the incubating soils did not differ between cover crop treatments, these data may not reflect C availability in the field since soils were air-dried and re-wetted for incubation, which changes C availability (Bijay-Singh et al., 1988). In incubating soils from N225 without cover crop, glucose additions had no effect on N₂O emissions, in contrast to other fertilized treatments (Figure 5). High NO₃ concentrations in soils from this treatment may have inhibited denitrification (Lalisse-Grundmann et al., 1988; Weier et al., 1993).

Since some N₂O produced was likely reduced to N₂ before diffusing from the soil, N₂O emission data did not represent total denitrification in incubating soils. Some studies have found decreased N₂O emissions with glucose additions to soils incubating at high water content, attributed to greater reduction of N₂O to N₂ with increased C availability (e.g. Miller

et al., 2008; Sánchez-Martín et al., 2008; Andersen and Petersen, 2009). In this experiment, substantial reduction of N_2O to N_2 was likely inhibited by the high NO_3 concentrations in the fertilized soils (Blackmer and Bremner, 1978). Because more electrons are accepted by N in the reduction of NO_3 to N_2O (4 electrons) than the reduction of N_2O to N_2 (1 electron), the mass of glucose added may not have provided sufficient demand for electron acceptors to decrease N_2O emissions. In addition to denitrification, nitrification could have produced N_2O in the incubating soils. However, substantial production of N_2O by nitrification was unlikely given the high WFPS at which soils were incubated (Firestone and Davidson, 1989).

Cover Crop Effects on N_2O Emissions

The use of cover crops has been proposed to reduce N_2O emissions from cropping systems, attributed to their ability to reduce mineral N availability for N_2O -producing processes (McSwiney et al., 2010; Eagle and Olander, 2012). However, the effects of cover crops on factors influencing N_2O emissions are complex and interact with climate and management practices. When N fertilizer is broadcast and incorporated, N immobilization by cover crops may reduce mineral N concentrations sufficiently to reduce N_2O emissions. In contrast, in this study, fertilizer was applied in concentrated bands, coincident with high rainfall and soil moisture (data not shown). Due to the high mineral N availability in the bands, N immobilization in cover crop residue may have been insufficient to decrease N_2O emissions (Jarecki et al., 2009). At the same time, given the favorable conditions for denitrification in the bands soon after N application, C substrate provided by cover crop residue likely stimulated N_2O emissions. Though not observed in this study, cover crops can also decrease

soil moisture by evapotranspiration (Thorup-Kristensen et al., 2003), which may decrease N₂O emissions in systems where N₂O emissions are limited by O₂ diffusion.

Given the complex effects of cover crops on the factors influencing N₂O emissions, and the interactions of these factors with climate and management practices, general expectations that cover crops decrease N₂O emissions from agricultural soils (e.g. Eagle and Olander, 2012) may be unfounded. However, N₂O emissions from the soil surface do not represent the total effect of cover crops on N₂O emissions or global warming potential of the cropping system. Non-legume cover crops, such as rye, can decrease NO₃ leaching losses to aquatic systems (Tonitto et al., 2006). Since models of agricultural N₂O emissions estimate a percentage of leached NO₃ to be emitted as N₂O downstream (Smith et al., 2007), cover crops may decrease total N₂O emissions by decreasing NO₃ leaching (Snyder et al., 2009). Cover crops also affect the soil C balance, both by providing an input of C and by affecting crop yield (Thorup-Kristensen et al., 2003; Eagle and Olander, 2012).

Conclusion

Despite reducing soil NO₃ concentrations at a range of N fertilizer rates, a rye cover crop preceding maize only decreased N₂O emissions in unfertilized treatments, but increased N₂O emissions at an N rate close to the economic optimum. The degree of mineral N immobilization by over-wintering cover crops may not be sufficient to reduce soil N₂O emissions, while the input of mineralizable C in cover crop residue following N application may stimulate N₂O emissions. The effects of cover crops on multiple factors controlling N₂O

emissions, including C availability, should be considered in evaluations of their net effect on N₂O emissions from agricultural soils.

Acknowledgements

The authors thank Fernando Míguez and Daniel Barker for assistance. This research is part of a regional collaborative project supported by the USDA-NIFA, “Cropping Systems Coordinated Agricultural Project (CAP): Climate Change, Mitigation, and Adaptation in Corn-based Cropping Systems.” Project Web site: sustainablecorn.org. The field project is supported in part by the Iowa Department of Agriculture and Land Stewardship, Division of Soil Conservation, through funds appropriated by the Iowa General Assembly.

References

- Adviento-Borbe MAA, Haddix ML, Binder DL, Walters DT, Dobermann A. 2007. Soil greenhouse gas fluxes and global warming potential in four high-yielding maize systems. *Global Change Biology* 13: 1972-1988.
- Andersen AJ, Petersen SO. 2009. Effects of C and N availability and soil-water potential interactions on N₂O evolution and PLFA composition. *Soil Biology and Biochemistry* 41: 1726-1733.
- Baggs EM, Watson CA, Rees RM. 2000. The fate of nitrogen from incorporated cover crop and green manure residues. *Nutrient Cycling in Agroecosystems* 56: 153-163.
- Bijay-Singh J, Ryden JC, Whitehead DC. 1988. Some relationships between denitrification potential and fractions of organic carbon in air-dried and field-moist soils. *Soil Biology and Biochemistry* 20: 737-741.
- Blackmer AM, Bremner JM. 1978. Inhibitory effect of nitrate on reduction of N₂O to N₂

- by soil microorganisms. *Soil Biology and Biochemistry* 10: 187-191.
- Bremner JM. 1997. Sources of nitrous oxide in soils. *Nutrient Cycling in Agroecosystems* 49: 7-16.
- Burford JR, Bremner JM. 1975. Relationships between the denitrification capacities of soils and total, water-soluble and readily decomposable soil organic matter. *Soil Biology and Biochemistry* 7: 389-394.
- Eagle AJ, Olander LP. 2012. Greenhouse gas mitigation with agricultural land management activities in the United States: a side-by-side comparison of biophysical potential. *Advances in Agronomy* 115: 79-179.
- Firestone MK, Davidson EA. 1989. Microbiological basis of NO and N₂O production and consumption in soil. In: Andreae, M.O., Schimel, D.S. (Eds.), *Exchange of Trace Gases between Terrestrial Ecosystems and the Atmosphere*. John Wiley & Sons, New York, pp. 7-21.
- Gillam KM, Zebarth BJ, Burton DL. 2008. Nitrous oxide emissions from denitrification and the partitioning of gaseous losses as affected by nitrate and carbon addition and soil aeration. *Canadian Journal of Soil Science* 88: 133-143.
- Hood-Nowotny R, Umana NH-N, Inselbacher E, Oswald-Lachouani P, Wanek W. 2010. Alternative methods for measuring inorganic, organic, and total dissolved nitrogen in soil. *Soil Science Society of America Journal* 74: 1018-1027.
- Iqbal J, Castellano MJ, Parkin TB. 2012. Evaluation of photoacoustic infrared spectroscopy for simultaneous measurement of N₂O and CO₂ concentrations and fluxes at the soil surface. *Global Change Biology* doi:10.1111/gcb.12021
- Iowa State University Agronomy Extension. 2012. Corn nitrogen rate calculator. Accessed September 2012. <http://extension.agron.iastate.edu/soilfertility/nrate.aspx>
- Jarecki MK, Parkin TB, Chan ASK, Kaspar TC, Moorman TB, Singer JW, Kerr BJ, Hatfield JL, Jones R. 2009. Cover crop effects on nitrous oxide emission from a manure-treated Mollisol. *Agriculture, Ecosystems and Environment* 134: 29-35.
- Lalisse-Grundmann G, Brunel B, Chalamet A. 1988. Denitrification in a cultivated soil:

- optimal glucose and nitrate concentrations. *Soil Biology and Biochemistry* 20: 839-844.
- McKenney DJ, Wang S-W, Drury CF, Findlay WI. 1995. Denitrification, immobilization, and mineralization in nitrate limited and nonlimited residue-amended soil. *Soil Science Society of America Journal* 59: 118-124.
- McSwiney CP, Robertson GP. 2005. Nonlinear response of N₂O flux to incremental fertilizer addition in a continuous maize (*Zea mays* L.) cropping system. *Global Change Biology* 11: 1712-1719.
- McSwiney CP, Snapp SS, Gentry LE. 2010. Use of N immobilization to tighten the N cycle in conventional agroecosystems. *Ecological Applications* 20: 648-662.
- Miller MN, Zebarth BJ, Dandie CE, Burton DL, Goyer C, Trevors JT. 2008. Crop residue influence on denitrification, N₂O emissions and denitrifier community abundance in soil. *Soil Biology and Biochemistry* 40: 2553-2562.
- Mosier AR, Halvorson AD, Reule CA, Liu XJ. 2006. Net global warming potential and greenhouse gas intensity in irrigated cropping systems in northeastern Colorado. *Journal of Environmental Quality* 35: 1584-1598.
- Myrold DD, Tiedje JM. 1985. Establishment of denitrification capacity in soil: effects of carbon, nitrate, and moisture. *Soil Biology and Biochemistry* 17: 819-822.
- Ostrom NE, Sutka R, Ostrom PH, Grandy AS, Huizinga KM, Gandhi H, von Fischer JC, Robertson GP. 2010. Isotopologue data reveal bacterial denitrification as the primary source of N₂O during a high flux event following cultivation of a native temperate grassland. *Soil Biology and Biochemistry* 42: 499-506.
- Parkin TB, Venterea RT. 2010. Chapter 3: Chamber-based trace gas flux measurements. In: Follett, R.F. (Ed.), *Sampling Protocols*. USDA-ARS, pp. 3-1 to 3-39. Available at: www.ars.usda.gov/research/GRACEnet
- Petersen SO, Mutegi JK, Hansen EM, Munkholm LJ. 2011. Tillage effects on N₂O emissions as influenced by a winter cover crop. *Soil Biology and Biochemistry* 43: 1509-1517.
- Reay DS, Davidson EA, Smith KE, Smith P, Melillo JM, Dentener F, Crutzen PJ.

2012. Global agriculture and nitrous oxide emissions. *Nature Climate Change* 2: 410-416.
- Robertson GP, Vitousek PM. 2009. Nitrogen in agriculture: balancing the cost of an essential resource. *Annual Review of Environment and Resources* 34: 97-125.
- Sainz Rozas HR, Echeverría HE, Picone LI. 2001. Denitrification in maize under no-tillage: effect of nitrogen rate and application time. *Soil Science Society of America Journal* 65: 1314-1323.
- Sánchez-Martín L, Vallejo A, Dick J, Skiba UM. 2008. The influence of soluble carbon and fertilizer nitrogen on nitric oxide and nitrous oxide from two contrasting agricultural soils. *Soil Biology and Biochemistry* 40: 142-151.
- Sarkodie-Addo J, Lee HC, Baggs EM. 2003. Nitrous oxide emissions after application of inorganic fertilizer and incorporation of green manure residues. *Soil Use and Management* 19: 331-339.
- Sawyer JE, Nafziger E, Randall G, Bundy L, Rehm G, Joern B. 2006. Concepts and rationale for regional nitrogen rate guidelines for corn. Publication PM2015, Iowa State University Extension, Ames, IA, USA.
- Sehy U, Ruser R, Munch JC. 2003. Nitrous oxide fluxes from maize fields: relationship to yield, site-specific fertilization, and soil conditions. *Agriculture, Ecosystems and Environment* 99: 97-111.
- Smith P, Martino D, Cai, Z, et al. 2007. Chapter 8: Agriculture. In: Metz B, Davidson OR, Bosch PR, Dave R, Meyer LA (Eds.), *Climate Change 2007: Mitigation. Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK, pp. 497-540.
- Smith DR, Hernandez-Ramirez G, Armstrong SD, Bucholtz DL, Stott DE. 2011. Fertilizer and tillage management impacts on non-carbon-dioxide greenhouse gas emissions. *Soil Science Society of America Journal* 75: 1070-1082.
- Snyder CS, Bruulsema TW, Jensen TL, Fixen PE. 2009. Review of greenhouse gas emissions from crop production systems and fertilizer management effects. *Agriculture, Ecosystems and Environment* 133: 247-266.

- Stehfest E, Bouwman L. 2006. N₂O and NO emission from agricultural fields and soils under natural vegetation: summarizing available measurement data and modeling of global annual emissions. *Nutrient Cycling in Agroecosystems* 74: 207-228.
- Thorup-Kristensen K, Magid J, Jensen LS. 2003. Catch crops and green manures as biological tools in nitrogen management in temperate zones. *Advances in Agronomy* 79: 227-302.
- Tiedje JM. 1994. Denitrifiers. In: Weaver, R.W., Angle, J.S., Bottomley, P.S. (Eds.), *Methods of Soil Analysis, Part 2 – Microbiological and Biochemical Properties*. Soil Science Society of America, Madison, WI, USA, pp. 245-268.
- Tonitto C, David MB, Drinkwater LE. 2006. Replacing bare fallows with cover crops in fertilizer-intensive cropping systems: a meta-analysis of crop yield and N dynamics. *Agriculture, Ecosystems and Environment* 112: 58-72.
- Van Groenigen JW, Kasper GJ, Velthof GL, van den Pol-van Dasselaar A, Kuikman PJ. 2004. Nitrous oxide emissions from silage maize fields under different mineral nitrogen fertilizer and slurry applications. *Plant and Soil* 263: 101-111.
- Weier KL, Doran JW, Power JF, Walters DT. 1993. Denitrification and the dinitrogen / nitrous oxide ratio as affected by soil water, available carbon, and nitrate. *Soil Science Society of America Journal* 57: 66-72.

Table 3.1. Maize grain yields (at 155 g kg⁻¹ moisture) at the field site harvested 5 Oct 2011.

N rate	No rye cover crop	With rye cover crop
kg N ha ⁻¹	----- Mg ha ⁻¹ -----	
0	5.18 (0.39) [†]	3.22 (0.33)
135	9.83 (0.28)	9.63 (0.44)
225	10.51 (0.59)	10.37 (0.86)

[†] Values in parentheses are standard deviations.

Table 3.2. Analysis of variance of cumulative N₂O produced over 10 days in incubating soils. Cumulative emission data were log-transformed before statistical analysis.

Treatment	<i>P</i> > <i>F</i>
Cover Crop	0.209
N Fertilizer Rate	0.062
Cover Crop × N Fertilizer Rate	0.136
NO ₃ Addition	0.608
Glucose Addition	0.071
Cover Crop × NO ₃ Addition	0.662
N Fertilizer Rate × NO ₃ Addition	0.008
Cover Crop × N Fertilizer Rate × NO ₃ Addition	0.677
Cover Crop × Glucose Addition	0.721
N Fertilizer Rate × Glucose Addition	0.140
Cover Crop × N Fertilizer Rate × Glucose Addition	0.004
NO ₃ Addition × Glucose Addition	0.167
Cover Crop × NO ₃ Addition × Glucose Addition	0.305
N Fertilizer Rate × NO ₃ Addition × Glucose Addition	0.136
Cover Crop × N Fertilizer Rate × NO ₃ Addition × Glucose Addition	0.512




Figure 3.1. Mean soil NO₃-N concentrations over the period of field gas flux measurements (11 Apr - 3 Oct 2011) with one-way analysis of variance results. Data were log-transformed for statistical analysis. Error bars show standard errors of means of 4 replicates.




Figure 3.2. Cumulative $\text{N}_2\text{O-N}$ emissions from the field study (11 Apr - 3 Oct 2011) with one-way analysis of variance results. Data were log-transformed for statistical analysis. Error bars show standard errors of means of 4 replicates.




Figure 3.3. Initial NO₃-N concentrations in incubating soils collected from the fertilizer bands of the field site 13-16 June 2011 with one-way analysis of variance results. Data were log-transformed for statistical analysis. Error bars show standard errors of means of 4 replicates.




Figure 3.4. Cumulative N₂O-N emissions over 10 days from soils collected from the fertilizer bands of the field site and incubated at 0.8 cm³ cm⁻³ WFPS, showing the effect of NO₃ addition at each N fertilizer rate across cover crop and glucose addition treatments. Error bars show standard errors of means of 4 replicates.




Figure 3.5. Cumulative $\text{N}_2\text{O-N}$ emissions over 10 days from soils collected from the fertilizer bands of the field site and incubated at $0.8 \text{ cm}^3 \text{ cm}^{-3}$ WFPS, showing the effect of glucose addition at each cover crop \times N fertilizer rate combination. Error bars show standard errors of means of 4 replicates.

CHAPTER 4. GENERAL CONCLUSIONS

Since N and C cycles are coupled in agroecosystems, management practices meant to reduce loss of N forms must take into account the central role of organic C inputs and cycling in agricultural soils. The studies presented in this thesis provide two examples of how soil C inputs influence loss of reactive N forms (NO_3 and N_2O). These studies combine insights from the fields of ecosystem ecology and soil biology and biochemistry and apply these insights to evaluate the impact of agricultural management practices on environmental quality.

In Chapter 2, denitrification was identified as the most important sink for NO_3 inputs into a perennial vegetation buffer (PVB) ecosystem. This result contrasted with the hypothesis, based on preliminary data and previous studies of ecosystem N retention, that soil organic matter would be the most important NO_3 sink. While the literature on NO_3 retention has focused on riparian buffer systems such as wetlands and riparian forests, the site of this study provides a unique opportunity to investigate the ability of PVBs integrated into row crop landscapes to decrease NO_3 losses to streams. Despite the differences in hydrology and ecology between this system and most riparian systems, denitrification has been identified as the principal NO_3 sink in both systems. Because $\text{NO}_3\text{-N}$ is lost from the ecosystem during denitrification, these results indicate that PVB will remain effective as NO_3 sinks in the long term. These results also indicate that organic C inputs from perennial vegetation are a key factor increasing denitrification in PVB soils compared to row crop soils, leading to decreased NO_3 leaching losses. Further research is needed to determine if these results would

be consistent across varying hydrological and geological conditions, as well as to quantify N₂O emissions produced by denitrification in these systems.

In Chapter 3, an overwintering non-legume cover crop was found to have inconsistent effects on N₂O emissions from the soil surface across a wide range of N fertilizer rates. This study provided an example of a case in which a cover crop increased N₂O emissions from fertilized soil, in contrast to previous studies which have proposed cover crops as a practice to decrease N₂O emissions. Furthermore, in this study, mineralizable C inputs from cover crop residue were identified as a likely factor influencing increased N₂O emissions from fertilized soils. These results agree with many studies showing limitation of N₂O emissions by mineralizable C availability in fertilized soils with high NO₃ availability. Further studies are needed to determine the full effect of cover crops on N₂O emissions to the atmosphere, since decreases in NO₃ leaching with a cover crop may lead to less N₂O emissions downstream. Nevertheless, this study adds to a body of evidence that cover crop effects on N₂O emissions from cropping system soils should not be expected to be consistent across management practices, climatic conditions, and other factors.

ACKNOWLEDGEMENTS

I would like to sincerely thank Michael Castellano for providing me with the opportunity to study at here at Iowa State, for giving me a tremendous amount of freedom, encouragement, guidance and support in my research, and for his patience with its completion. I thank my committee members Matthew Helmers, Kirsten Hofmockel, and John Sawyer for the time and attention they dedicated to my research here. I thank Ehsan Razavy Toosi for providing much enjoyable company and enlightening conversation during our two years together at ISU; he has been missed as a good friend and mentor to many here, myself included. Many thanks to Javed Iqbal and Kim Brown for their support, assistance, and patience as lab mates and colleagues. I would like to thank Carolina Córdova for her assistance and for the good company, conversation, and inspiration she has so generously provided. In addition to the co-authors and others acknowledged in the chapters, I would like to thank the following individuals for their assistance and contributions to this research: Charl Lombard, Shonda Eister, Samantha Lodge, Danielle Koester, Dan Anderson, Whitney Bouma, Marlín Pérez Suárez, Sarah Hargreaves, Ryan Williams, and Andrea Basche. Finally, I would like to thank my Mom and Dad, Amma and Papa, and Noah and Shannon for their tremendous family love and support.