

1-2013

Soybean Response to Drought

Mark A. Licht

Iowa State University, lichtma@iastate.edu

David Wright

Iowa State University

Andrew W. Lenssen

Iowa State University, alenssen@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/extension_ag_pubs



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Agronomy and Crop Sciences Commons](#)

Recommended Citation

Licht, Mark A.; Wright, David; and Lenssen, Andrew W., "Soybean Response to Drought" (2013). *Agriculture and Environment Extension Publications*. 190.

http://lib.dr.iastate.edu/extension_ag_pubs/190

Iowa State University Extension and Outreach publications in the Iowa State University Digital Repository are made available for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current publications and information from Iowa State University Extension and Outreach, please visit <http://www.extension.iastate.edu>.

Soybean Response to Drought

Severe drought is a serious, albeit infrequent, stressor affecting Iowa soybean production. Severe statewide drought in Iowa has occurred in 1956, 1988, and 2012. However, soil water deficits caused by brief periods of low rainfall occur more frequently and can occur more than once during the growing season. During these periods soybean can experience drought-like conditions when available soil water becomes short. Soybean appears to be somewhat resilient to brief periods of drought, or water deficit, but the timing, duration, and severity of the stress plays a major role in determining the impact on yield. Soybean is most susceptible to water deficits during germination and the reproductive growth stages.

The soybean seed needs to imbibe 50 percent of its weight in water to germinate and begin to elongate the radicle and hypocotyl. Once plant growth begins, plant tissues consist of about 90 percent water. Water plays an important role in plant developmental and respiratory systems, helps regulate leaf temperature, and carries nutrients and sugars to key metabolic systems.

Soybean can respond to a water deficit as early as two days after germination (20). Laboratory experiments showed that hypocotyl elongation was progressively inhibited as available soil water decreased while root elongation was unaffected. As plants age, the response to a water deficit is similar. In fact, there have been several reports that show continued root growth after soil water deficits slow or stop shoot growth in several crops (7,10,15,18). This response to drought stress enables the roots to



Soybean wilting in response to drought.

grow deeper into the soil profile to access additional available soil water.

Short term, moderate water deficits during vegetative growth stages generally do not impact soybean yield. Shoot growth rate reduction during short-term drought can be followed by compensatory growth during periods of rainfall (5). However, severe and/or long-term water deficits can cause irreversible dehydration of vegetative tissue resulting in low grain yield or plant death.

Soybean yield is most sensitive to water deficits during the pod filling stage of development (19). A reduction in pod number, as much as 20 percent, as a result of flower abortion, is often reported as being highly affected by soil water deficits (1,17). Seeds per pod and seed size are also impacted (2) but to a lesser extent than pod number. Stressed plants often mature earlier, shortening the grain filling period causing reduced seed weight and yield (17).

ROOT RESPONSE TO WATER DEFICITS

More than 76 percent of soybean roots grow in the top 16 inches of soil with more than 50 percent growing in the top eight inches (6). Soybean rooting depth has been measured at 4.9 to 6.5 feet under normal field conditions (9,12,20). Under ideal conditions, soybean roots can grow downward at a rate of 0.5 inches per day from planting to R1 (beginning bloom) (3) and 2.3 inches per day from R1 to R3 (beginning pod) (9). Downward growth continues through growth stages R5 to R6 (seed fill) under normal field conditions. However, downward growth after R5 appears to be variety specific (9).

Soybean roots in the upper soil profile grow slowly under water deficit conditions. Soybean compensates by partitioning photosynthates to roots to enable them to grow more rapidly in the lower soil profile (4,12). The increased root growth rate appears to coincide with water-deficits encountered three to five days previously (6). Growth rates of water stressed plants return to normal when soil water levels return to normal.

KEY POINTS

- Moderate water deficits during vegetative growth stages generally do not impact yield.
- Soybean yield loss from water deficits is caused by a reduction in flower number, pod number, and seed fill. Yield loss can also occur from reduced N fixation by nodules.
- Management practices that leave low amounts of residue on the soil surface and increase water runoff can increase soil water deficits.
- Adopting production practices that reduce soil compaction and promote increased water infiltration rate can improve yield when drought conditions occur.

IMPACT ON NITROGEN FIXATION

Water deficits reduce symbiotic nitrogen fixation of soybean roots. Biological N fixation accounts for 25 to 75 percent of the annual N uptake for the soybean crop (21) and is essential to producing high yielding soybeans. A decline in nitrogen fixation causes yield reduction by restricting nitrogen (N) used for protein production, which is critical for seed fill. The decrease in N fixation during water deficits is associated with limited water (11) and oxygen supply to the nodules, a decrease in rhizobia (13), and nodule carbon shortage (8). Oxygen limitation is not the sole factor in reducing nitrogen fixation and seems to be involved only in the initial stages of water-deficit related stress in decreasing nodule activity (14).

SUMMARY

Soybean appears to be somewhat resilient to brief periods of drought, or water deficit, but the timing, duration, and severity of the stress plays a major role in determining the impact on yield. Soybean is most susceptible to water deficits during germination and the reproductive growth stages.

Soybean yield loss from water deficits is caused by a reduction in flower number, pod number, and seed fill during the reproductive stages of growth. Yield loss can also occur from reduced N fixation by nodules.

Management practices that leave high amounts of residue on the soil surface and improve soil structure can increase water infiltration rate and increase soybean yield. Adopting production practices that reduce soil compaction also can improve soybean yield when drought conditions occur by improving root growth and nodule function.

REFERENCES

1. Cox, W.J., and G.D. Jolliff. 1986. Growth and yield of sunflower and soybean under soil water deficits. *Agronomy Journal* 78:226-230.
2. Desclaux, D., T. Huynh, and P. Roumet. 2000. Identification of soybean plant characteristics that indicate the timing of drought stress. *Crop Science* 40:716-722.
3. Fehr, W.R., and C.E. Caviness. 1977. Stages of soybean development. Spec. Rep. 80. Iowa Agric. Home Econ. Exp. Stn., Iowa State Univ., Ames.
4. Garay, A.F., and W.W. Wilhlem. 1983. Root system characteristics of two soybean isolines undergoing water stress conditions. *Agronomy Journal* 75:973-977.
5. Hoogenboom, G., C.M. Peterson, and M.G. Huck. 1987a. Shoot growth rate of soybean as affected by drought stress. *Agronomy Journal* 79:598-607.
6. Hoogenboom, G., M.G. Huck, and C.M. Peterson. 1987b. Root growth rate of soybean as affected by drought stress. *Agronomy Journal* 79:607-614.
7. Hsiao, T.C., and J. Jing. 1987. Leaf and root expansive growth in response to water deficits. In D.J. Cosgrove, D.P. Knievel, eds, *Physiology of cell expansion during plant growth*. American Society of Plant Physiologists, Rockville, MD. pp 180-192.
8. King, C.A. and L.C. Purcell. 2001. Soybean nodule size and relationship to nitrogen fixation response to water deficit. *Crop Science* 41:1099-1107.
9. Kaspar, T.C., C.D. Stanley, and H.M. Taylor. 1978. Soybean root growth during the reproductive stages of development. *Agronomy Journal* 70:1105-1107.
10. Kramer, P.J. 1983. *Water relations of plants*. Academic Press, New York.
11. Ladrera, R.D. Marino, E. Larrainzar, E.M. Gonzalez and C. Arrese-Igor. 2007. Reduced carbon availability to bacteroids and elevated ureides in nodules, but not in shoots, are involved in the nitrogen fixation response to early drought in soybean. *Plant Physiology* 145:539-546
12. Mayaki, W.C., I.D. Teare, and L.R. Stone. 1976. Top and root growth of irrigated and nonirrigated soybeans. *Crop Science* 16:92-94.
13. Peña-Cabriaes, J.J. and M. Alexandar. 1979. Survival of rhizobium in soils undergoing drying. *Soil Science Society of America Journal* 43:962-966.
14. Serraj, R. 2003. Effects of drought stress on legume symbiotic nitrogen fixation: physiological mechanisms. *Indian Journal of Experimental Botany* 10:1136-1141.
15. Sharp, R.E. and W.J. Davies. 1979. Solute regulation and growth by roots and shoots of water-stressed maize plants. *Planta* 147:43-49.
16. Sinclair, T.R., C.D. Messina, A. Beatty, M. Samples. 2010. Assessment across the United States of the Benefits of Altered Soybean Drought Traits. *Agronomy Journal* 102:475-482.
17. Sionit, N. and P.J. Kramer. 1977. Effect of water stress during different stages of growth of soybean. *Agronomy Journal* 69:274-278.
18. Westgate, M.E. and J.S. Boyer. 1985. Osmotic adjustment and the inhibition of leaf root, stem and silk growth at low-water potentials in maize. *Planta* 164-540-549.
19. Westgate, M.E., and C.M. Peterson. 1993. Flower and pod development in water-deficient soybean (*Glycine max* L. Merr.). *Journal Experimental Botany* 44:109-117.
20. Willett, S.T. and H.M. Taylor, 1978. Water uptake of soybean roots as affected by their depth and by soil water content. *Journal Agricultural Science* 90:205-213.
21. Zapata, F., S.K.A. Danso, G. Hardarson, and M. Fried. 1987. Time course of nitrogen fixation in field-grown soybean using nitrogen-15 methodology. *Agronomy Journal* 79:172-176.

Prepared by Mark Licht, Iowa State University Extension and Outreach, and David Wright and Andy Lenssen, Department of Agronomy, Iowa State University.

... and justice for all

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Many materials can be made available in alternative formats for ADA clients. To file a complaint of discrimination, write USDA, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914 in cooperation with the U.S. Department of Agriculture. Cathann A. Kress, director, Cooperative Extension and Outreach, Iowa State University of Science and Technology, Ames, Iowa.