

1-2013

Humic and Fulvic Acids and Their Potential in Crop Production

David Wright
Iowa State University

Andrew W. Lenssen
Iowa State University, alenssen@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/extension_ag_pubs

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Agronomy and Crop Sciences Commons](#)

Recommended Citation

Wright, David and Lenssen, Andrew W., "Humic and Fulvic Acids and Their Potential in Crop Production" (2013). *Agriculture and Environment Extension Publications*. 187.
http://lib.dr.iastate.edu/extension_ag_pubs/187

Iowa State University Extension and Outreach publications in the Iowa State University Digital Repository are made available for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current publications and information from Iowa State University Extension and Outreach, please visit <http://www.extension.iastate.edu>.

Humic and Fulvic Acids and Their Potential in Crop Production

The use of soilborne compounds to protect seeds and enhance seedling growth is being investigated globally. Humic and fulvic acids are commonly used in organic crop production, greenhouse cropping of vegetables, and in horticultural programs. They also may have uses in Iowa's large-scale production systems to improve seedling health and plant response to stress.

Humic compounds such as humic acid and fulvic acid have been shown to stimulate plant growth in terms of increasing plant height and dry or fresh weight as well as enhancing nutrient uptake. These effects seem to depend on the concentration (2,3,5) and source of the substance and on the plant species.

Research is being conducted at Iowa State University to evaluate if humic and fulvic acid based compounds influence soybean yield and quality. Because these compounds are believed to enhance root development, their influence on soybean grown in the presence of the soybean cyst nematode (SCN) is also being investigated. SCN accounts for an estimated \$1.5 billion in crop loss annually.

Humic and fulvic acids are the most characteristic compounds of soil humic substances. Humic substances are formed through the microbial degradation of plant material and the brown to black substances are the primary constituents of soil organic matter.



Photo: Wayne Fredericks

Humic substances have the ability to hold seven times their volume in water, a greater water holding capacity than clay soils. Water stored within the topsoil enables plant roots to quickly access available nutrients required for plant growth and yield.

Despite the amount of research information on humic and fulvic acids, there is surprisingly little on their effects on soybean and corn.

Early research on soybean found a significant increase in root dry matter after the roots and shoots were exposed to humic acid and fulvic acid. Nodule weight was also increased but the number of nodules was inversely related to increasing amounts of humic acid (5).

Stimulation of root and shoot growth by humic acid was also reported in corn (2). They reported an increase in branching and root hair development of corn roots when plants were

grown in a nutrient solution containing humic acids. Similar results were reported on tobacco roots (3).

Root proliferation is a benefit from applications of humic and fulvic acids at low concentrations. These stimulatory effects also have been directly correlated with enhanced uptake of nitrogen, phosphorus, sulfur, zinc, and iron (4).

However, use of these compounds at high concentrations also has been shown to decrease root and shoot growth (2,3,4,5). The source of the humic acid substances and the concentrations used in treatments are not uniform among experiments so few conclusions can be drawn from these studies. Hartwigsen and Evans (1) reported an increase in root fresh weight of horticultural crops when 2500 ppm humic acid was used while Lee and Bartlett (2) reported the optimum concentration for corn was 5 ppm humic acid.

CONCLUSIONS

Many positive effects of humic substances such as humic and fulvic acids on plant growth and development have been reported but their effects appear to be dependent on several chemical and soil factors. In leguminous crops such as soybean, humic acid applications reduced nodule number but remaining nodules were larger in size. At low concentrations, humic acid increased vegetative growth of soybean and corn in laboratory trials.

Prepared by David Wright and Andy Lensen, Department of Agronomy, Iowa State University.

... and justice for all

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Many materials can be made available in alternative formats for ADA clients. To file a complaint of discrimination, write USDA, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Cathann A. Kress, director, Cooperative Extension and Outreach, Iowa State University of Science and Technology, Ames, Iowa.

Understanding humic substances and the way they influence plant growth and development and nutrient uptake is crucial to developing sustainable cropping systems that improve overall soil quality. Although humic substances are currently widely used in organic production systems, they may have uses in Iowa's large-scale production systems to improve seedling health. Stimulation of root growth may improve plant resistance to disease, and plant response to feeding by herbivores and nematodes, and water stress caused by drought.

REFERENCES

1. Hartwigsen, J.A. and M.R. Evans. 2000. Humic acid seed and substrate treatments promote seedling root development. *HortScience*, 35(7):1231-1233.
2. Lee, Y.S. and R.J. Bartlett. 1976. Stimulation of plant growth by humic substances. *Soil Science Society of America Journal*, 40:876-879.
3. Mylonas, V.A. and C.B. McCants. 1980a. Effects of humic and fulvic acids on growth of tobacco. *Plant and Soil*, 54: 485-490.
4. Mylonas, V.A. and C.B. McCants. 1980b. Effects of humic and fulvic acids on growth of tobacco 2. Tobacco growth and ion uptake. *Journal of Plant Nutrition*, 2: 377-393.
5. Tan, K.H. and D. Tantiwiranond. 1983. Effect of humic acids on nodulation and dry matter production of soybean peanut, and clover. *Soil Science Society of America Journal*, 47:1121-1124.